



Metabolically engineered cells for the production of resveratrol or an oligomeric or glycosidically-bound derivative thereof

Katz, Michael; Smits, Hans Peter; Förster, Jochen; Bredal Nielsen, Jens

Publication date:
2006

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Katz, M., Smits, H. P., Förster, J., & Bredal Nielsen, J. (2006). Metabolically engineered cells for the production of resveratrol or an oligomeric or glycosidically-bound derivative thereof. (Patent No. WO2006089898).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 August 2006 (31.08.2006)

PCT

(10) International Publication Number
WO 2006/089898 A1

(51) International Patent Classification:

C12N 9/00 (2006.01) *C12N 9/88* (2006.01)
C12N 9/02 (2006.01) *C12P 7/22* (2006.01)
C12N 9/10 (2006.01)

(21) International Application Number:

PCT/EP2006/060154

(22) International Filing Date:

21 February 2006 (21.02.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0503657.9 22 February 2005 (22.02.2005) GB

(71) Applicant (for all designated States except US): **FLUX-OME SCIENCES A/S** [DK/DK]; Diplomvej 378, DK-2800 Kgs. Lyngby (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KATZ, Michael** [SE/SE]; Ormvråksgatan 9, S-215 62 Malmö (SE). **SMITS, Hans Peter** [NL/DK]; Søllerødvej 39B, DK-2840 Holte (DK). **FÖRSTER, Jochen** [DE/DK]; Halmtorvet 7, 5tv, DK-1700 Copenhagen (DK). **NIELSEN, Jens Bredal** [DK/DK]; Hans Bruunsvej 8, DK-2920 Charlottenlund (DK).

(74) Agent: **SMART, Peter, John**; Fulwood House, 12 Fulwood Place, London, London GB WC1V 6HR (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METABOLICALLY ENGINEERED CELLS FOR THE PRODUCTION OF RESVERATROL OR AN OLIGOMERIC OR GLYCOSIDICALLY-BOUND DERIVATIVE THEREOF

(57) Abstract: A recombinant micro-organism producing resveratrol by a pathway in which phenylalanine ammonia lyase (PAL) produces trans-cinnamic acid from phenylalanine, cinnamate 4- hydroxylase (C4H) produces 4-coumaric acid from said trans- cinnamic acid, 4-coumarate-CoA ligase (4CL) produces 4- coumaroyl CoA from said 4-coumaric acid, and resveratrol synthase (VST) produces said resveratrol from said 4- coumaroyl CoA, or in which L-phenylalanine- or tyrosine- ammonia lyase (PAL/TAL) produces 4-coumaric acid, 4- coumarate-CoA ligase (4CL) produces 4-coumaroyl CoA from said 4-coumaric acid, and resveratrol synthase (VST) produces said resveratrol from said 4-coumaroyl CoA. The micro-organism may be a yeast, fungus or bacterium including *Saccharomyces cerevisiae*, *E. coli*, *Lactococcus lactis*, *Aspergillus niger*, or *Aspergillus oryzae*.



WO 2006/089898 A1

**METABOLICALLY ENGINEERED CELLS FOR THE PRODUCTION OF
RESVERATROL OR AN OLIGOMERIC OR GLYCOSIDICALLY-BOUND
DERIVATIVE THEREOF.**

FIELD OF THE INVENTION

5

This invention relates generally to the production of the polyphenol resveratrol or an oligomeric or glycosidically bound derivative thereof such as its β -glucoside piceid using microbial cells. Furthermore, it relates to the use of
10 naturally occurring or recombinant micro-organisms that produce resveratrol or such a derivative for production of food, feed and beverages.

BACKGROUND OF THE INVENTION

15

Production of chemicals from micro-organisms has been an important application of biotechnology. Typically, the steps in developing such a bio-production method may include 1) selection of a proper micro-organism host, 2) elimination of
20 metabolic pathways leading to by-products, 3) deregulation of desired pathways at both enzyme activity level and the transcriptional level, and 4) overexpression of appropriate enzymes in the desired pathways. In preferred aspect, the present invention has employed combinations of the steps
25 above to redirect carbon flow from phenylalanine or tyrosine through enzymes of the plant phenylpropanoid pathway which supplies the necessary precursor for the desired biosynthesis of resveratrol.

Resveratrol (or 3,4,5-trihydroxystilbene) is a phytophenol belonging to the group of stilbene phytoalexins, which are low-molecular-mass secondary metabolites that constitute the active defence mechanism in plants in response to infections or other stress-related events. Stilbene phytoalexins contain the stilbene skeleton (*trans*-1,2-diphenylethylene) as their common basic structure: that may be supplemented by addition of other groups as well (Hart and Shrimpton, 1979, Hart, 1981). Stilbenes have been found in certain trees (angio-sperms, gymnosperms), but also in some herbaceous plants (in species of the *Myrtaceae*, *Vitaceae* and *Leguminosae* families). Said compounds are toxic to pests, especially to fungi, bacteria and insects. Only few plants have the ability to synthesize stilbenes, or to produce them in an amount that provides them sufficient resistance to pests.

The synthesis of the basic stilbene skeleton is pursued by stilbene synthases. So far, two enzymes have been designated as a stilbene synthase; pinosylvine synthase and resveratrol synthase. To date, the groundnut (*Arachis hypogaea*) resveratrol synthase has been characterised in most detail, such that most of the properties are known (Schoppner and Kindl, 1984). Substrates that are used by stilbene synthases are malonyl-CoA, cinnamoyl-CoA or coumaroyl-CoA. These substances occur in every plant because they are used in the biosynthesis of other important plant constituents as well such as flavonoids, flower pigments and lipids.

Resveratrol (Fig. 1 *trans*-form) consists of two closely connected phenol rings and belongs therefore to the polyphenols. While present in other plants, such as eucalyptus, spruce, and lily, and in other foods such as

mulberries and peanuts, resveratrol's most abundant natural sources are *Vitis vinifera*, *-labrusca*, and *-muscadine (rotundifolia)* grapes, which are used to make wines. The compound occurs in the vines, roots, seeds, and stalks, but
5 its highest concentration is in the skin (Celotti et al., 1996), which contains 50-100 µg/g. (Jang et al. 1997). During red wine vinification the grape skins are included in the must, in contrast to white wine vinification, and therefore resveratrol is found in small quantities in red
10 wine only. Resveratrol has, besides its antifungal properties, been recognized for its cardioprotective- and cancer chemopreventive activities; it acts as a phytoestrogen, an inhibitor of platelet aggregation (Kopp et al, 1998; Gehm et al 1997; Lobo et al 1995), and an
15 antioxidant (Jang et al., 1997; Huang 1997). These properties explain the so-called French Paradox, i.e. the wine-drinking French have a low incidence of coronary heart disease despite a low-exercise, high-fat diet. Recently it has been shown that resveratrol can also activate the SIR2
20 gene in yeast and the analogous human gene SIRT1, which both play a key role in extending life span. Ever since, attention is very much focused on the life-span extending properties of resveratrol (Hall, 2003, Couzin, 2004).

25 American health associations, such as the Life Extension Foundation, are promoting the vast beneficial effects of this drug, and thereby propelling the ideal conditions for a successful commercialisation. Present production processes rely mostly upon extraction of resveratrol, either from the
30 skin of grape berries, or from Knot weed. This is a labour intensive process and generates low yield which, therefore, prompts an incentive for the development of novel, more efficient and high-yielding production processes.

In plants, the phenylpropanoid pathway is responsible for the synthesis of a wide variety of secondary metabolic compounds, including lignins, salicylates, coumarins, hydroxycinnamic amides, pigments, flavonoids and phytoalexins. Indeed formation of resveratrol in plants proceeds through the phenylpropanoid pathway. The amino acid L-phenylalanine is converted into *trans*-cinnamic acid through the non-oxidative deamination by L-phenylalanine ammonia lyase (PAL) (Fig 2). Next, *trans*-cinnamic acid is hydroxylated at the *para*-position to 4-coumaric acid (4-hydroxycinnamic acid) by cinnamate-4-hydroxylase (C4H), a cytochrome P450 monooxygenase enzyme, in conjunction with NADPH:cytochrome P450 reductase (CPR). The 4-coumaric acid, is subsequently activated to 4-coumaroyl-CoA by the action of 4-coumarate-CoA ligase (4CL). Finally, resveratrol synthase (VST) catalyses the condensation of a phenylpropane unit of 4-coumaroyl-CoA with malonyl CoA, resulting in formation of resveratrol.

20

Recently, a yeast was disclosed that could produce resveratrol from 4-coumaric acid that is found in small quantities in grape must (Becker et al. 2003). The production of 4-coumaroyl-CoA, and concomitant resveratrol, in laboratory strains of *S. cerevisiae*, was achieved by co-expressing a heterologous coenzyme-A ligase gene, from hybrid poplar, together with the grapevine resveratrol synthase gene (*vst1*). The other substrate for resveratrol synthase, malonyl-CoA, is already endogenously produced in yeast and is involved in *de novo* fatty-acid biosynthesis. The study showed that cells of *S. cerevisiae* could produce minute amounts of resveratrol, either in the free form or in

30

the glucoside-bound form, when cultured in synthetic media that was supplemented with 4-coumaric acid.

However, said yeast would not be suitable for a commercial application because it suffers from low resveratrol yield, and requires addition of 4-coumaric acid, which is only present in few industrial media. In order to facilitate and broaden the application of resveratrol as both a pharmaceutical and nutraceutical, it is therefore highly desirable to obtain a yeast that can produce resveratrol directly from glucose, without addition of 4-coumaric acid.

A recent study (Ro and Douglas, 2004) describes the reconstitution of the entry point of the phenylpropanoid pathway in *S. cerevisiae* by introducing PAL, C4H and CPR from Poplar. The purpose was to evaluate whether multienzyme complexes (MECs) containing PAL and C4H are functionally important at this entry point into phenylpropanoid metabolism. By feeding the recombinant yeast with [3H]-phenylalanine it was found that the majority of metabolized [3H]-phenylalanine was incorporated into 4-[3H]-coumaric acid, and that phenylalanine metabolism was highly reduced by inhibiting C4H activity. Moreover, PAL-alone expressers metabolized very little phenylalanine into cinnamic acid. When feeding [3H]-phenylalanine and [14C]-*trans*-cinnamic acid simultaneously to the triple expressers, no evidence was found for channeling of the endogenously synthesized [3H]-*trans*-cinnamic acid into 4-coumaric acid. Therefore, efficient carbon flux from phenylalanine to 4-coumaric acid via reactions catalyzed by PAL and C4H does not appear to require channeling through a MEC in yeast, and sheer biochemical coupling of PAL and C4H seems to be sufficient to drive carbon flux into the phenylpropanoid pathway. In

yet another study (Hwang *et al.*, 2003) production of plant-specific flavanones by *Escherichia coli* was achieved through expression of an artificial gene cluster that contained three genes of a phenyl propanoid pathway of various
5 heterologous origins; PAL from the yeast *Rhodotorula rubra*, 4CL from the actinomycete *Streptomyces coelicolor*, and chalcone synthase (CHS) from the licorice plant *Glycyrrhiza echinata*. These pathways bypassed C4H, because the bacterial 4CL enzyme ligated coenzyme A to both *trans*-cinnamic acid
10 and 4-coumaric acid. In addition, the PAL from *Rhodotorula rubra* uses both phenylalanine and tyrosine as the substrates. Therefore, *E. coli* cells containing the gene clusters and grown on glucose, produced small amounts of two flavanones, pinocembrin (0.29 g/l) from phenylalanine and
15 naringenin (0.17 g/l) from tyrosine. In addition, large amounts of their precursors, 4-coumaric acid and *trans*-cinnamic acid (0.47 and 1.23 mg/liter respectively), were accumulated. Moreover, the yields of these compounds could be increased by addition of phenylalanine and tyrosine.

20

Whereas the enzyme from dicotyledonous plants utilizes only phenylalanine efficiently, several studies indicated that PAL from monocotyledonous plants, and some micro-organisms, utilizes tyrosine as well (Rösler *et al.*, 1997). In such
25 reactions the enzyme activity is designated tyrosine ammonia lyase (TAL, figure 3). Conversion of tyrosine by TAL results in the direct formation of 4-coumaric acid without the intermediacy of C4H and CPR. Mostly both activities reside on the same polypeptide and have very similar catalytic
30 efficiencies, in spite of large differences in K_m and turnover number. However, most PAL/TAL enzymes from plants prefer phenylalanine rather than tyrosine. The level of TAL activity is mostly lower than PAL activity, but the

magnitude of this difference varies over a wide range. For example, the parsley enzyme has a K_m for phenylalanine of 15-25 μM and for tyrosine 2.0-8.0 mM with turnover numbers 22 s^{-1} and 0.3 s^{-1} respectively. In contrast, the maize enzyme
5 has a K_m for phenylalanine only 15-fold higher than for tyrosine, and turnover numbers about 10-fold higher. Moreover, in the red yeasts, *Rhodotorula glutinis* (*Rhodospiridium toruloides*) and *-rubra*, the TAL catalytic activity is close to the PAL catalytic activity with a ratio
10 of TAL/PAL of approximately 0.58. It is believed that the PAL enzyme in these yeasts degrades phenylalanine as a catabolic function and the *trans*-cinnamic acid formed is converted to benzoate and other cellular materials, whereas in plants it is thought to be merely a regulatory enzyme in
15 the biosynthesis of lignin, isoflavonoids and other phenylpropanoids.

Recently, an open reading frame was found in the bacterium *Rhodobacter capsulatus* that encodes a hypothetical
20 biosynthetic tyrosine ammonia lyase (TAL) that is involved in the biosynthesis of the chromophore of the photoactive yellow protein (Kyndt et al., 2002). This was the first time that a PAL-homologous gene was found in bacteria. The TAL gene was isolated and overproduced in *Escherichia coli*. The
25 K_m and k_{cat} values for the conversion of tyrosine to 4-coumaric acid were 15.6 μM and 27.7 s^{-1} respectively, and for conversion of L-phenylalanine to *trans*-cinnamic acid were 1277 μM and 15.1 s^{-1} respectively. As a consequence of the smaller K_m and a slightly larger k_{cat} , the enzyme shows a
30 strong preference for tyrosine over L-phenylalanine, with a catalytic efficiency (K_m/k_{cat}) for tyrosine of approximately 150-fold larger than for phenylalanine. The kinetic studies established that tyrosine, and not L-phenylalanine, is the

natural substrate of the enzyme under physiological conditions. Very recently a study described the heterologous coexpression of phenylalanine ammonia lyase, cinnamate-4-hydroxylase, 4-coumarate-CoA-ligase and chalcone synthase, for the production of flavonoids in *E. coli* (Watts et al., 2004). The simultaneous expression of all four genes, however, was not successful because of a nonfunctional cinnamate-4-hydroxylase. The substitution of phenylalanine ammonia lyase and cinnamate-4-hydroxylase by a new tyrosine ammonia lyase that was cloned from *Rhodobacter sphaeroides*, could, however, solved the problem and led to high-level production of the flavonone naringenin. Furthermore, said tyrosine ammonia lyase from *Rhodobacter sphaeroides* is also used for heterologous production of 4-coumaric acid (i.e. para-hydroxycinnamic acid) in *Escherichia coli* (US-A-2004059103). Evenmore, further methods for development of a biocatalyst for conversion of glucose into 4-coumaric acid are described. US-A-2004023357 discloses a tyrosine ammonia lyase from the yeast *Trichosporon cutaneum* for the production of coumaric acid in *Escherichia coli* and *Saccharomyces cerevisiae*. US-A-2001053847 describes the incorporation of the wild type PAL from the yeast *Rhodotorula glutinis* into *E. coli*, underlining the ability of the wildtype PAL to convert tyrosine directly to 4-coumaric acid. Moreover, there is also exemplification of incorporation of the wildtype PAL from the yeast *Rhodotorula glutinis*, plus a plant C4H and CPR into *E. coli* and *S. cerevisiae*. Also described is the development of a biocatalyst through mutagenesis of the wild type yeast PAL *Rhodotorula glutinis* with enhanced TAL activity (US-A-6521748). Neither of the aforementioned patents claim the incorporation of 4CL and VST for the production of resveratrol.

Recently, evidence was shown that the filamentous fungi *A. oryzae* contained the enzyme chalcone synthase (CHS) that is normally involved in the biosynthesis of flavonoids, such as naringenin, in plants (Seshime *et al.*, 2005). Indeed it was also shown that *A. oryzae* contained the major set of genes responsible for phenylpropanoid-flavonoid metabolism, i.e. PAL, C4H and 4CL. However, there is no evidence that *A. oryzae* contained a stilbene synthase such as resveratrol synthase.

The present invention now provides a micro-organism having an operative metabolic pathway comprising at least one enzyme activity, said pathway producing 4-coumaric acid and producing resveratrol therefrom or an oligomeric or glycosidically-bound derivative thereof. Such a micro-organism may be naturally occurring and may be isolated by suitable screening procedures, but more preferably is genetically engineered.

Preferably, said resveratrol or derivative is produced in a reaction catalysed by an enzyme in which endogenous malonyl-CoA is a substrate, and preferably said resveratrol is produced from 4-coumaroyl-CoA.

Said resveratrol or derivative is preferably produced from 4-coumaroyl-CoA by a resveratrol synthase which is preferably expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.

Generally herein, unless the context implies otherwise, references to resveratrol include reference to oligomeric or

glycosidically bound derivatives thereof, including particularly piceid.

Thus, in certain preferred embodiments, said resveratrol
5 synthase is a resveratrol synthase (EC 2.3.1.95) from a plant belonging to the genus of *Arachis*, e.g. *A. glabatra*, *A. hypogaea*, a plant belonging to the genus of *Rheum*, e.g. *R. tataricum*, a plant belonging to the genus of *Vitus*, e.g. *V. labrusca*, *V. riparia*, *V. vinifera*, or any one of the
10 genera *Pinus*, *Picea*, *Lilium*, *Eucalyptus*, *Parthenocissus*, *Cissus*, *Calochortus*, *Polygonum*, *Gnetum*, *Artocarpus*, *Nothofagus*, *Phoenix*, *Festuca*, *Carex*, *Veratrum*, *Bauhinia* or *Pterolobium*.

15 Preferably, said 4-coumaric acid is produced from *trans*-cinnamic acid, suitably by an enzyme in a reaction catalysed by said enzyme in which oxygen is a substrate, NADH or NADPH is a cofactor and NAD^+ or NADP^+ is a product.

20 Thus, said 4-coumaric acid may be produced from *trans*-cinnamic acid by a cinnamate 4-hydroxylase, which preferably is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.

25 In certain preferred embodiments, including those referred to in the paragraphs above, said cinnamate-4-hydroxylase is a cinnamate-4-hydroxylase (EC 1.14.13.11) from a plant or a micro-organism. The plant may belong to the genus of *Arabidopsis*, e.g. *A. thaliana*, a plant belonging to the
30 genus of *Citrus*, e.g. *C. sinensis*, *C. x paradisi*, a plant belonging to the genus of *Phaseolus*, e.g. *P. vulgaris*, a plant belonging to the genus of *Pinus*, e.g. *P. taeda*, a plant belonging to the genus of *Populus*, e.g. *P. deltoides*,

P. tremuloides, *P. trichocarpa*, a plant belonging to the genus of *Solanum*, e.g. *S. tuberosum*, a plant belonging to the genus of *Vitus*, e.g. *Vitus vinifera*, a plant belonging to the genus of *Zea*, e.g. *Z. mays*, or other plant genera e.g. *Ammi*, *Avicennia*, *Camellia*, *Camptotheca*, *Catharanthus*,
5 *Glycine*, *Helianthus*, *Lotus*, *Mesembryanthemum*, *Physcomitrella*, *Ruta*, *Saccharum*, *Vigna*. The micro-organism might be a fungus belonging to the genus *Aspergillus*, e.g. *A. oryzae*.

10

Preferably, said 4-coumaric acid is produced from tyrosine in a reaction catalysed by an enzyme in which ammonia is produced and suitably, said 4-coumaric acid is produced from tyrosine by a L-phenylalanine ammonia lyase or a tyrosine
15 ammonia lyase, e.g. tyrosine ammonia lyase (EC 4.3.1.5) from yeast or bacteria. Suitably, the tyrosine ammonia lyase is from the yeast *Rhodotorula rubra* or from the bacterium *Rhodobacter capsulatus*.

20 Optionally, said tyrosine ammonia lyase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.

Alternatively, said *trans*-cinnamic acid may be produced from
25 L-phenylalanine in a reaction catalysed by an enzyme in which ammonia is produced and suitably said *trans*-cinnamic acid is formed from L-phenylalanine by a phenylalanine ammonia lyase.

30 In certain preferred embodiments, said L-phenylalanine ammonia lyase is a L-phenylalanine ammonia lyase (EC 4.3.1.5) from a plant or a micro-organism. The plant may belong to the genus of *Arabidopsis*, e.g. *A. thaliana*, a

plant belonging to the genus of *Brassica*, e.g. *B. napus*, *B. rapa*, a plant belonging to the genus of *Citrus*, e.g. *C. reticulata*, *C. clementinus*, *C. limon*, a plant belonging to the genus of *Phaseolus*, e.g. *P. coccineus*, *P. vulgaris*, a plant belonging to the genus of *Pinus*, e.g. *P. banksiana*, *P. monticola*, *P. pinaster*, *P. sylvestris*, *P. taeda*, a plant belonging to the genus of *Populus*, e.g. *P. balsamifera*, *P. deltoides*, *P. Canadensis*, *P. kitakamiensis*, *P. tremuloides*, a plant belonging to the genus of *Solanum*, e.g. *S. tuberosum*, a plant belonging to the genus of *Prunus*, e.g. *P. avium*, *P. persica*, a plant belonging to the genus of *Vitus*, e.g. *Vitus vinifera*, a plant belonging to the genus of *Zea*, e.g. *Z. mays* or other plant genera e.g. *Agastache*, *Ananas*, *Asparagus*, *Bromheadia*, *Bambusa*, *Beta*, *Betula*, *Cucumis*, *Camellia*, *Capsicum*, *Cassia*, *Catharanthus*, *Cicer*, *Citrullus*, *Coffea*, *Cucurbita*, *Cynodon*, *Daucus*, *Dendrobium*, *Dianthus*, *Digitalis*, *Dioscorea*, *Eucalyptus*, *Gallus*, *Ginkgo*, *Glycine*, *Hordeum*, *Helianthus*, *Ipomoea*, *Lactuca*, *Lithospermum*, *Lotus*, *Lycopersicon*, *Medicago*, *Malus*, *Manihot*, *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Olea*, *Oryza*, *Pisum*, *Persea*, *Petroselinum*, *Phalaenopsis*, *Phyllostachys*, *Physcomitrella*, *Picea*, *Pyrus*, *Quercus*, *Raphanus*, *Rehmannia*, *Rubus*, *Sorghum*, *Sphenostylis*, *Stellaria*, *Stylosanthes*, *Triticum*, *Trifolium*, *Triticum*, *Vaccinium*, *Vigna*, *Zinnia*. The micro-organism might be a fungus belonging to the genus *Agaricus*, e.g. *A. bisporus*, a fungus belonging to the genus *Aspergillus*, e.g. *A. oryzae*, *A. nidulans*, *A. fumigatus*, a fungus belonging to the genus *Ustilago*, e.g. *U. maydis*, a bacterium belonging to the genus *Rhodobacter*, e.g. *R. capsulatus*, a yeast belonging to the genus *Rhodotorula*, e.g. *R. rubra*.

Suitably, said L-phenylalanine ammonia lyase is expressed in said micro-organism from nucleic acid coding for said enzyme

which is not native to the micro-organism.

Preferably, 4-coumaroyl-CoA is formed in a reaction catalysed by an enzyme in which ATP and CoA are substrates and ADP is a product and suitably 4-coumaroyl-CoA is formed
5 in a reaction catalysed by a 4-coumarate-CoA ligase.

Said 4-coumarate-CoA ligase may be a 4-coumarate-CoA ligase (EC 6.2.1.12) from a plant, a micro-organism or a nematode.
10 The plant may belong to the genus of *Abies*, e.g. *A. beshanzuensis*, *B. firma*, *B. holophylla*, a plant belonging to the genus of *Arabidopsis*, e.g. *A. thaliana*, a plant belonging to the genus of *Brassica*, e.g. *B. napus*, *B. rapa*, *B. oleracea*, a plant belonging to the genus of *Citrus*, e.g.
15 *C. sinensis*, a plant belonging to the genus of *Larix*, e.g. *L. decidua*, *L. gmelinii*, *L. griffithiana*, *L. himalaica*, *L. kaempferi*, *L. laricina*, *L. mastersiana*, *L. occidentalis*, *L. potaninii*, *L. sibirica*, *L. speciosa*, a plant belonging to the genus of *Phaseolus*, e.g. *P. acutifolius*, *P. coccineus*, a
20 plant belonging to the genus of *Pinus*, e.g. *P. armandii*, *P. banksiana*, *P. pinaster*, a plant belonging to the genus of *Populus*, e.g. *P. balsamifera*, *P. tomentosa*, *P. tremuloides*, a plant belonging to the genus of *Solanum*, e.g. *S. tuberosum*, a plant belonging to the genus of *Vitus*, e.g.
25 *Vitus vinifera*, a plant belonging to the genus of *Zea*, e.g. *Z. mays*, or other plant genera e.g. *Agastache*, *Amorpha*, *Cathaya*, *Cedrus*, *Crocus*, *Festuca*, *Glycine*, *Juglans*, *Keteleeria*, *Lithospermum*, *Lolium*, *Lotus*, *Lycopersicon*, *Malus*, *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Nothotsuga*,
30 *Oryza*, *Pelargonium*, *Petroselinum*, *Physcomitrella*, *Picea*, *Prunus*, *Pseudolarix*, *Pseudotsuga*, *Rosa*, *Rubus*, *Ryza*, *Saccharum*, *Suaeda*, *Thellungiella*, *Triticum*, *Tsuga*. The micro-organism might be a filamentous fungi belonging to the

genus *Aspergillus*, e.g. *A. flavus*, *A. nidulans*, *A. oryzae*,
A. fumigatus, a filamentous fungus belonging to the genus
Neurospora, e.g. *N. crassa*, a fungus belonging to the genus
Yarrowia, e.g. *Y. lipolytica*, a fungus belonging to the
5 genus of *Mycosphaerella*, e.g. *M. graminicola*, a bacterium
belonging to the genus of *Mycobacterium*, e.g. *M. bovis*, *M.*
leprae, *M. tuberculosis*, a bacterium belonging to the genus
of *Neisseria*, e.g. *N. meningitidis*, a bacterium belonging to
the genus of *Streptomyces*, e.g. *S. coelicolor*, a bacterium
10 belonging to the genus of *Rhodobacter*, e.g. *R. capsulatus*, a
nematode belonging to the genus *Ancylostoma*, e.g. *A.*
ceylanicum, a nematode belonging to the genus
Caenorhabditis, e.g. *C. elegans*, a nematode belonging to the
genus *Haemonchus*, e.g. *H. contortus*, a nematode belonging to
15 the genus *Lumbricus*, e.g. *L. rubellus*, a nematode belonging
to the genus *Meilodogyne*, e.g. *M. hapla*, a nematode
belonging to the genus *Strongyloidus*, e.g. *S. ratti*, *S.*
stercoralis, a nematode belonging to the genus *Pristionchus*,
e.g. *P. pacificus*.

20

Optionally, a NADPH:cytochrome P450 reductase (CPR) has been
recombinantly introduced into said micro-organism. This may
be a plant CPR introduced into a non-plant micro-organism.
Alternatively, a native NADPH:cytochrome P450 reductase
25 (CPR) has been overexpressed in said micro-organism.

In certain preferred embodiments, including those referred
to in the paragraphs above, said NADPH:cytochrome P450
reductase is a NADPH:cytochrome P450 reductase (EC 1.6.2.4)
30 from a plant belonging to the genus of *Arabidopsis*, e.g. *A.*
thaliana, a plant belonging to the genus of *Citrus*, e.g. *C.*
sinensis, *C. x paradisi*, a plant belonging to the genus of
Phaseolus, e.g. *P. vulgaris*, a plant belonging to the genus

of *Pinus*, e.g. *P. taeda*, a plant belonging to the genus of *Populus*, e.g. *P. deltoides*, *P. tremuloides*, *P. trichocarpa*, a plant belonging to the genus of *Solanum*, e.g. *S. tuberosum*, a plant belonging to the genus of *Vitus*, e.g. *Vitus vinifera*, a plant belonging to the genus of *Zea*, e.g. *Z. mays*, or other plant genera e.g. *Ammi*, *Avicennia*, *Camellia*, *Camptotheca*, *Catharanthus*, *Glycine*, *Helianthus*, *Lotus*, *Mesembryanthemum*, *Physcomitrella*, *Ruta*, *Saccharum*, *Vigna*.

10

Whilst the micro-organism may be naturally occurring, preferably at least one copy of at least one genetic sequence encoding a respective enzyme in said metabolic pathway has been recombinantly introduced into said micro-organism.

15

Additionally or alternatively to introducing coding sequences coding for a said enzyme, one may provide one or more expression signals, such as promoter sequences, not natively associated with said coding sequence in said organism. Thus, optionally, at least one copy of a genetic sequence encoding a tyrosine ammonia lyase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism, and/or at least one copy of a genetic sequence encoding a L-phenylalanine ammonia lyase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

20

25

Optionally, at least one copy of a genetic sequence encoding cinnamate 4-hydroxylase, whether native or not, is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

30

Optionally, at least one copy of a genetic sequence encoding a 4-coumarate-CoA ligase, whether native or not, is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

Optionally, at least one copy of a genetic sequence encoding a resveratrol synthase, whether native or not, is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

Expression signals include nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Such sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

20

In certain aspects the invention provides a metabolically engineered micro-organism having an operative metabolic pathway in which a first metabolite is transformed into a second metabolite in a reaction catalysed by a first enzyme, said reaction step producing ammonia, and in which said second metabolite is transformed into a third metabolite in a reaction catalysed by a second enzyme, in which oxygen is a substrate, NADPH or NADH is a cofactor and NADP⁺ or NAD⁺ is a product, and in which said third metabolite is transformed into a fourth metabolite in a reaction catalysed by a third enzyme in which ATP and CoA is a substrate, and ADP is a product, and in which said fourth metabolite is transformed

into a fifth metabolite in a reaction catalysed by a fourth enzyme in which endogenous malonyl-CoA is a substrate.

The present invention also provides a metabolically engineered micro-organism having an operative metabolic pathway in which a first metabolite is transformed into a said third metabolite catalysed by a first enzyme, said reaction step producing ammonia, without the involvement of said second enzyme, and in which said third metabolite is transformed into a said fourth metabolite in a reaction catalysed by a said third enzyme in which ATP and CoA is a substrate, and ADP is a product, and in which said fourth metabolite is transformed into a said fifth metabolite in a reaction catalysed by a said fourth enzyme in which endogenous malonyl-CoA is a substrate.

The micro-organisms described above include ones containing one or more copies of an heterologous DNA sequence encoding phenylalanine ammonia lyase operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding cinnamate-4-hydroxylase operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding 4-coumarate-CoA-ligase operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding resveratrol synthase operatively associated with an expression signal.

They include also ones lacking cinnamate-4-hydroxylase activity, and containing one or more copies of a heterologous DNA sequence encoding tyrosine ammonia lyase operatively associated with an expression signal, and

containing one or more copies of an heterologous DNA sequence encoding 4-coumarate-CoA-ligase operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding
5 resveratrol synthase operatively associated with an expression signal.

In the present context the term "micro-organism" relates to microscopic organisms, including bacteria, microscopic
10 fungi, including yeast.

More specifically, the micro-organism may be a fungus, and more specifically a filamentous fungus belonging to the genus of *Aspergillus*, e.g. *A. niger*, *A. awamori*, *A. oryzae*,
15 *A. nidulans*, a yeast belonging to the genus of *Saccharomyces*, e.g. *S. cerevisiae*, *S. kluyveri*, *S. bayanus*, *S. exiguus*, *S. sevazzi*, *S. uvarum*, a yeast belonging to the genus *Kluyveromyces*, e.g. *K. lactis* *K. marxianus* var. *marxianus*, *K. thermotolerans*, a yeast belonging to the genus
20 *Candida*, e.g. *C. utilis* *C. tropicalis*, *C. albicans*, *C. lipolytica*, *C. versatilis*, a yeast belonging to the genus *Pichia*, e.g. *P. stipidis*, *P. pastoris*, *P. sorbitophila*, or other yeast genera, e.g. *Cryptococcus*, *Debaromyces*, *Hansenula*, *Pichia*, *Yarrowia*, *Zygosaccharomyces* or
25 *Schizosaccharomyces*. Concerning other micro-organisms a non-exhaustive list of suitable filamentous fungi is supplied: a species belonging to the genus *Penicillium*, *Rhizopus*, *Fusarium*, *Fusidium*, *Gibberella*, *Mucor*, *Mortierella*, *Trichoderma*.

30

Concerning bacteria a non-exhaustive list of suitable bacteria is given as follows: a species belonging to the genus *Bacillus*, a species belonging to the genus

Escherichia, a species belonging to the genus *Lactobacillus*,
a species belonging to the genus *Lactococcus*, a species
belonging to the genus *Corynebacterium*, a species belonging
to the genus *Acetobacter*, a species belonging to the genus
5 *Acinetobacter*, a species belonging to the genus *Pseudomonas*,
etc.

The preferred micro-organisms of the invention may be *S.*
cerevisiae, *A. niger*, *A. oryzae*, *E. coli*, *L. lactis* or *B.*
10 *subtilis*.

The constructed and engineered micro-organism can be
cultivated using commonly known processes, including
chemostat, batch, fed-batch cultivations, etc.

15 The constructed and engineered micro-organism can be
cultivated using commonly known processes, including
chemostat, batch, fed-batch cultivations, etc.

Thus, the invention includes a method for producing
resveratrol or an oligomeric or glycosidically-bound
derivative thereof comprising contacting a non-plant cell
with a carbon substrate in the substantial absence of an
20 external source of 4-coumaric acid, said cell having the
capacity to produce resveratrol or an oligomeric or
glycosidically-bound derivative thereof under the
conditions, in which the micro-organism may be selected from
the group consisting of fungi and bacteria, especially
25 yeast.

Said carbon substrate is optionally selected from the group
of fermentable carbon substrates consisting of
monosaccharides, oligosaccharides and polysaccharides, e.g.
30 glucose, fructose, galactose, xylose, arabinose, mannose,
sucrose, lactose, erythrose, threose, and/or ribose. Said
carbon substrate may additionally or alternatively be
selected from the group of non-fermentable carbon substrates

including ethanol, acetate, glycerol, and/or lactate. Said non-fermentable carbon substrate may additionally or alternatively be selected from the group of amino acids and may be phenylalanine and/or tyrosine.

5

In an alternative aspect, the invention includes a method for producing resveratrol or an oligomeric or glycosidically-bound derivative thereof through heterologous expression of nucleotide sequences encoding phenylalanine ammonia lyase, cinnamate 4-hydroxylase, 4-coumarate-CoA ligase and resveratrol synthase and also a method for producing resveratrol through heterologous expression of nucleotide sequences encoding tyrosine ammonia lyase, 4-coumarate-CoA ligase and resveratrol synthase.

15

Resveratrol or an oligomeric or glycosidically-bound derivative thereof so produced may be used as a nutraceutical in a dairy product or a beverage such as beer.

Resveratrol produced according to the invention may be *cis*-resveratrol or *trans*-resveratrol, but it is to be expected that the *trans*- form will normally predominate.

20

BRIEF DESCRIPTION OF THE DRAWINGS

To assist in the ready understanding of the above description of the invention reference has been made to the accompanying drawings in which:

25

Figure 1 shows the chemical structure of *trans*-resveratrol;

Figure 2 shows the phenylpropanoid pathway utilising phenylalanine ammonia lyase acting on L-phenylalanine; and

Figure 3 shows the alternative pathway utilising
5 phenylalanine ammonia lyase acting on L-tyrosine.

Figure 4 shows the HPLC-chromatograms of extracts of *S. cerevisiae* strains FSSC-PALC4H4CLVST, FSSC-TAL4CLVST, grown on 100 g/l galactose. A chromatogram of 60 nanogram of pure
10 resveratrol is included.

Figure 5 shows the UV absorption spectrum for pure *trans*-resveratrol and *trans*-resveratrol produced by *S. cerevisiae* strain FSSC-PALC4H4CLVST, grown on 100 g/l galactose.
15

Figure 6 shows the HPLC-chromatograms of extracts from *E. coli* strains FSEC-TAL4CLVST and FSEC-control, grown on 50 g/l glucose.

Figure 7 shows the HPLC-chromatograms of extracts from *E. coli* strains FSEC-TAL4CLVST and FSEC-control, grown on 50 g/l glucose with addition of 20 mg/l coumaric acid. The UV absorption spectrum for *trans*-resveratrol produced in strain FSEC-TAL4CLVST is included.
20

25

The invention will be further described and illustrated by the following non-limiting examples.

EXAMPLES

Example 1

Isolation of genes encoding PAL, TAL, C4H, CPR, 4CL, and VST

5 Phenylalanine ammonia lyase (PAL2) (Cochrane et al., 2004; SEQ ID NO: 1, 2), cinnamate 4-hydroxylase (C4H) (Mizutani et al., 1997; SEQ ID NO: 3, 4) and 4-coumarate:CoenzymeA ligase (4CL1) (Hamberger and Hahlbrock 2004; Ehlting et al., 1999; SEQ ID NO: 5, 6) were isolated via PCR from *A. thaliana* cDNA
10 (BioCat, Heidelberg, Germany) using the primers in table 1. PAL2 and 4CL1 were chosen amongst several *A. thaliana* homologues due to favourable kinetic parameters towards cinnamic acid and coumaroyl-CoA, respectively (Cochrane et al., 2004; Hamberger and Hahlbrock 2004; Ehlting et al.,
15 1999).

The coding sequence of resveratrol synthase (VST) from *Rheum tataricum* (Samappito et al., 2003; SEQ ID NO: 7, 8) and tyrosine ammonia lyase (TAL) from *Rhodobacter capsulatus* (Kyndt et al., 2002; SEQ ID NO: 11, 12) were codon optimized
20 for expression in *S. cerevisiae* using the online service backtranslation tool at www.entelechon.com, yielding sequence SEQ ID NO: 9, 10 and SEQ ID NO: 13, 14 respectively. Oligos for the synthetic gene assembly were constructed at MWG Biotech and the synthetic gene was
25 assembled by PCR using a slightly modified method protocol of from Martin et al. (2003) described below.

Table 1. Primers and restriction sites for the amplification of genes			
Primer for amplification of gene* (Restriction sites are underlined)	Gene	Restriction site: primer	Restriction site: vector
5'-CGGAATTCTCATGGATCAAATCGAAGCAATGTT	PAL2	EcoR1	EcoR1
5'-CGACTAGTTTGTAGCAAATCGGAATCGGAGC	PAL2	Spe1	Spe1
5'-CGCTCGAGAT ATGGACCTCCTCTTGCTGGA	C4H	Xho1	Xho1
5'-CGGGTACCTTAACAGTTCCTTGCTTTCATAAC	C4H	Kpn1	Kpn1
5'-GCTCTAGACCT ATGGCGCCACAAGAACAAGCAGTTT	4CL1	Xba1	Spe1
5'-GCGGATCCCCT TCACAATCCATTTGCTAGTTT TGCC	4CL1	BamH1	BglII
5'-CC GGATCCAAATGGCCCCAGAAGAGAGCAGG	VST	BamH1	BamH1
5'-CG CTCGAGTTAAGTGATCAATGGAACCGAAGACAG	VST	Xho1	Xho1
5'-CCGAATTCCCATGACCCTGCAATCTCAAACAGCTAAAG	TAL	EcoR1	EcoR1
5'-CCACTAGTTTAAAGCAGGTGGATCGGCAGCT	TAL	Spe1	Spe1
5'-CCCTCGAGATCATGCCGTTTGAATAGACAACACCGA	CPR1	Xho1	Xho1
5'-CCAAGCTTATCGGGCTGATTACCAGACATCTTCTTG	CPR1	HindIII	HindIII
5'-CCGGATCCCCATGTCCTCTTCTTCTTCTTCGTCAAC	AR2	BamH1	BamH1
5'-CCCTCGAGGTGAGTGTGTGGCTTCAATAGTTT CG	AR2	Xho1	Xho1

* SEQ ID Nos 19-32

Primers from MWG for the assembly of the synthetic gene were dissolved in milliQ-water to a concentration of 100 pmole/ μ l. An aliquot of 5 μ l of each primer was combined in a totalmix and then diluted 10-fold with milliQ water. The gene was assembled via PCR using 5 μ l diluted totalmix per 50 μ l as template for fusion DNA polymerase (Finnzymes). The PCR programme was as follows: Initial 98 °C for 30 s., and then 30 cycles with 98 °C for 10 s., 40 °C for 1 min. and 72 °C at 1 min./1000 basepairs, and a final 72 °C for 5 min. From the resulting PCR reaction, 20 μ l was purified on 1% agarose gel. The result was a PCR smear and the regions around the wanted size were cut out from agarose gel and purified using the QiaQuick Gel Extraction Kit (Qiagen). A final PCR with the outer primers (for TAL and VST) in table 1 rendered the required TAL and VST genes. Point mutations were corrected using either the Quickchange site directed

mutagenesis II kit (Stratagene, La Jolla, CA), or using PCR from overlapping error free DNA stretches from several different *E. coli* subclones.

5 NADPH:Cytochrome P450 reductase (CPR) from *A. thaliana* (AR2) (Mizutani and Ohta, 1998; SEQ ID NO: 17, 18) and from *S. cerevisiae* (CPR1) (Aoyama *et al.*, 1978; SEQ ID NO: 15, 16), were isolated from *A. thaliana* cDNA (BioCat, Heidelberg, Germany) and *S. cerevisiae* genomic DNA, respectively, using the primers in table 1.

10

Example 2

Construction of a yeast vector for expression of PAL

15 The gene encoding PAL, isolated as described in example 1, was reamplified by PCR using forward- and reverse primers, with 5' overhangs containing EcoR1 and Spe1 restriction sites (table 1). The amplified PAL PCR product was digested with EcoR1/Spe1 and ligated into EcoR1/Spe1 digested pESC-URA vector (Stratagene), resulting in vector pESC-URA-PAL.
20 The sequence of the gene was verified by sequencing of two different clones.

.

Example 3

25 *Construction of a yeast vector for expression of PAL and C4H*

The gene encoding C4H, isolated as described in example 1, was amplified by PCR using the forward- and reverse primers, with 5' overhangs containing Xho1 and Kpn1 restriction
30 sites. The amplified C4H PCR-product was digested with Xho1/Kpn1 and ligated into similarly digested pESC-URA-PAL vector. The resulting plasmid, pESC-URA-PAL-C4H, contained

the genes encoding PAL and C4H under the control of the divergent GAL1/GAL10 promoter. The sequence of the gene encoding C4H was verified by sequencing of two different clones.

5

Example 4

Construction of a yeast vector for expression of 4CL

The gene encoding 4CL was isolated as described in example 1. The amplified 4CL PCR-product was digested with Xba1/BamH1 and ligated into Spe1/BglII digested pESC-TRP vector (Stratagene), resulting in vector pESC-TRP-4CL. Two different clones of pESC-TRP-4CL were sequenced to verify the sequence of the cloned gene.

15

Example 5

Construction of a yeast vector for expression of 4CL and VST

The gene encoding VST was isolated as described in example 1. The amplified synthetic VST gene was digested with BamH1/Xho1 and ligated into BamH1/Xho1 digested pESC-TRP-4CL (example 4). The resulting plasmid, pESC-TRP-4CL-VST, contained the genes encoding 4CL and VST under the control of the divergent GAL1/GAL10 promoter. The sequence of the gene encoding VST was verified by sequencing of two different clones of pESC-TRP-4CL-VST.

25

Example 6

Construction of a yeast vector for expression of TAL

30

The gene encoding TAL was isolated as described in example 1. The amplified synthetic TAL gene was digested with

EcoR1/Spe1 and ligated into EcoR1/Spe1-digested pESC-URA vector. The resulting plasmid, pESC-URA-TAL, contained the gene encoding for TAL under the control of the divergent GAL1/GAL10 promoter. The sequence was verified by sequencing
5 of two different clones of pESC-URA-TAL.

Example 7

Construction of a yeast vector for overexpression of S. cerevisiae endogenous CPR

10

The gene encoding CPR from *S. cerevisiae* (CPR1) was isolated as described in example 1. The amplified CPR1 gene was digested with Xho1/HindIII and ligated into Xho1/HindIII-digested pESC-LEU vector (Stratagene), resulting in vector
15 pESC-LEU-CPR1. The sequence was verified by sequencing of two different clones of pESC-LEU-CPR1.

Example 8

Construction of a yeast vector for overexpression of A. thaliana CPR (AR2)

20

The gene encoding CPR from *A. thaliana* (AR2) was isolated as described in example 1. The amplified AR2 gene was digested with BamH1/Xho1 and ligated into BamH1/Xho1 digested pESC-
25 LEU vector (Stratagene), resulting in vector pESC-LEU-AR2. The sequence was verified by sequencing of two different clones of pESC-LEU-AR2.

Example 9

Expression of the pathway to resveratrol in the yeast S. cerevisiae using PAL, C4H, 4CL and VST

5

Yeast strains containing the appropriate genetic markers were transformed with the vectors described in examples 2, 3, 4, 5, 6, 7 and 8, separately or in combination. The transformation of the yeast cell was conducted in accordance with methods known in the art, for instance, by using competent cells or by electroporation (see, e.g., Sambrook et al., 1989). Transformants were selected on medium lacking uracil and/or tryptophan and streak purified on the same medium.

15

S. cerevisiae strain CEN.PK 113-5D (MATa ura3) was transformed separately with the vector pESC-URA-PAL (example 2), yielding the strain FSSC-PAL, and with pESC-URA-PAL-C4H (example 3), resulting in the strain FSSC-PALC4H. *S. cerevisiae* strain FS01267 (MATa trp1 ura3) was co-transformed with pESC-URA-PAL-C4H and pESC-TRP-4CL (example 4), and the transformed strain was named FSSC-PALC4H4CL. The same strain was also co-transformed with pESC-URA-PAL-C4H and pESC-TRP-4CL-VST (example 5), resulting in the strain FSSC-PALC4H4CLVST.

25

Example 10

Expression of the pathway to resveratrol in S. cerevisiae using TAL, 4CL and VST.

30

S. cerevisiae strain CEN.PK 113-5D (MATa ura3) was transformed separately with the vector pESC-URA-TAL (example

6), yielding the strain FSSC-TAL. *S. cerevisiae* strain FS01267 (MATa trp1 ura3) was co-transformed with pESC-URA-TAL (example 6) and pESC-TRP-4CL (example 4), and the transformed strain was named FSSC-TAL4CL. The same strain
5 was also co-transformed with pESC-URA-TAL and pESC-TRP-4CL-VST (example 5), resulting in the strain FSSC-TAL4CLVST. Transformants were selected on medium lacking uracil and or tryptophan and streak purified on the same medium.

10 Example 11

Expression of the pathway to resveratrol in S.cerevisiae with overexpressed endogenous CPR.

S. cerevisiae strain FS01277 (MATa ura3 leu2 trp1) was co-
15 transformed with vectors pESC-URA-PAL-C4H (example 3), pESC-TRP-4CL (example 4), and pESC-LEU-CPR1 (example 7). The transformed strain was named FSSC-PALC4H4CLVSTCPR. Transformants were selected on medium lacking uracil and/or tryptophan and streak purified on the same medium.

20

Example 12

Expression of the pathway to resveratrol in S.cerevisiae with overexpressed A. thaliana CPR (AR2).

25 *S. cerevisiae* strain FS01277 (MATa ura3 leu2 trp1) was co-transformed with vectors pESC-URA-PAL-C4H (example 3), pESC-TRP-4CL (example 4), and pESC-LEU-AR2 (example 8). The transformed strain was named FSSC-PALC4H4CLVSTAR2. Transformants were selected on medium lacking uracil and or
30 tryptophan and streak purified on the same medium.

Example 13

Fermentation with recombinant yeast strains in shake flasks

5 The recombinant yeast strains were inoculated from agar plates with a sterile inoculation loop and grown in 200 ml defined mineral medium (Verduyn et al, 1992) that contained vitamins, trace elements, 5 g/l glucose and 40 g/l or 100 g/l galactose. The 500 ml stoppered shake flasks were incubated for three days at 30 °C and 160 rpm.

10

Example 14

Extraction of resveratrol

15 Cells were harvested by centrifugation 5000 g for 5 minutes. An aliquot of 50 ml of supernatant was extracted once with 20 ml ethyl acetate. The ethyl acetate was freeze dried and the dry product redissolved in 0.7 ml methanol and filtered into HPLC vials.

20 The cell pellet from 200 ml medium was dissolved in 1 to 2 ml water and divided into 3 fastprep tubes and broken with glass beads. The crude extracts from the three tubes were pooled into 10 ml 100 % methanol in a 50 ml sartorius tube and extracted on a rotary chamber for 48 hours in a dark cold room at 4 °C. After 48 hours the cell debris was removed
25 via centrifugation for 5 min. at 5000 g and the methanol was removed by freeze-drying overnight. The dried residue was redissolved in 1 ml phosphate-citrate buffer pH 5.4 and 10 units beta-glucosidase from almonds was added (Sigma) to release resveratrol from putatively glucoside-bound forms.
30 The mixture was incubated for three hours at 37 °C and then extracted twice with 1 ml ethyl acetate. The combined ethyl

acetate was freeze dried and the dry residue was redissolved in 0.7 ml methanol and filtered into HPLC vials.

Example 15

5 *Analysis of resveratrol*

Thin layer chromatography

10 A method based upon thin layer chromatography that enabled the quick separation of cinnamic, coumaric and resveratrol on the same TLC-plate was developed for quick screening analysis. An aliquot of 1 ml culture containing both cells and supernatant were extracted with 500 microliter ethyl acetate and centrifuged for 30 s. at 13000 rpm with a
15 microcentrifuge. The ethyl acetate was dried and redissolved in methanol. The extracts were analyzed on Silica G plates (0.2 mm Alugram SIL G/UV₂₅₄, Macherey-Nagel) containing a fluorescent indicator. The mobile phase was a mixture of chloroform, ethyl acetate and formic acid (25:10:1).

20

HPLC

For quantitative analysis of cinnamic acid, coumaric acid, and resveratrol, samples were subjected to separation by
25 high-performance liquid chromatography (HPLC) Agilent Series 1100 system (Hewlett Packard) prior to uv-diode-array detection at $\lambda = 306$ nm. A Phenomenex (Torrance, CA, USA) Luna 3 micrometer C18 (100 X 2.00 mm) column was used at 40 °C. As mobile phase a gradient of acetonitrile and milliq
30 water (both containing 50 ppm trifluoroacetic acid) was used at a flow of 0.4 ml/min. The gradient profile was linear from 15 % acetonitrile to 100 % acetonitrile over 20 min.

The elution times were approximately 3.4 min. for coumaric acid, 5.5 min. for free *trans*-resveratrol and 6.8 min. for cinnamic acid.

Pure resveratrol standard was purchased from Cayman chemical
5 company, whereas pure coumaric acid and cinnamic acid standards were purchased from and Sigma.

Results

Strains FSSC-PALC4H4CLVST and FSSC-TAL4CLVST, were cultivated
10 on 100 g/l galactose as described in example 13, and analyzed for their content of intracellular resveratrol according to example 14 and 15. Additionally, a control strain FSSC-control was included that contained the empty vectors pESC-URA and pESC-TRP only. The HPLC-analysis showed
15 that strains FSSC-PALC4H4CLVST and FSSC-TAL4CLVST contained a component with a retention time of 5.5 min. that was identical to *trans*-resveratrol (figure 4). Said result was confirmed by the UV absorption spectra that were similar to the absorption spectrum of pure *trans*-resveratrol (figure 5)
20 as well, with a λ_{\max} of approximately 306 nm.

The results, therefore, demonstrated the presence of an active phenyl-propanoid pathway in *S. cerevisiae* that led to *in vivo* production of *trans*-resveratrol. The production of resveratrol can most likely be improved by cultivating the
25 strains under well-defined growth conditions in batch- and continuous cultures, and/or optimizing the expression/activities of the individual enzymes.

Example 16

Construction of a bacterial vector for expression of TAL in Escherichia coli.

- 5 The gene encoding TAL, isolated as described in Example 1, was reamplified by PCR from the plasmid pESC-URA-TAL (example 6) using the forward primer 5'-CCG CTCGAG CGG ATG ACC CTG CAA TCT CAA ACA GCT AAA G-3' SEQ ID NO 33 and the reverse primer 5'-GC GGATCC TTA AGC AGG TGG ATC GGC AGC T-3' SEQ ID NO 34 with 5' overhangs containing the restriction sites XhoI and BamHI, respectively. The introduction of restriction sites at the 5' and 3' ends of the gene allowed ligation of the restricted PCR product into a pET16b vector (Novagen), digested with XhoI and BamHI to yield pET16b-TAL.
- 10
- 15 The pET16b vector contained both the ampicillin resistance gene, and the T7 promoter. Hence, above procedure resulted in a vector with an antibiotic selection marker that contained the gene encoding for TAL under the control of the T7 promoter. The sequence of the gene encoding TAL was
- 20 verified by sequencing of one clone of pET16b-TAL.

Example 17

Construction of a bacterial vector for expression of 4CL and VST in Escherichia coli.

- 25
- The gene encoding VST, isolated as described in example 1, was cut out with the restriction enzymes BamHI and XhoI from the digested plasmid pESC-TRP-4CL-VST (example 5), which contains the genes encoding 4CL and VST. The VST gene was
- 30 ligated into a pET26b vector (Novagen), containing the kanamycin resistance gene, digested with BamHI and SalI to yield pET26b-VST. The restriction enzymes XhoI and SalI have

compatible ends, which enabled proper ligation. The pET26b vector contained both the kanamycin resistance gene, and the T7 promoter. Hence, above procedure resulted in a vector with an antibiotic selection marker that contained the gene encoding for VST under the control of the T7 promoter.

The gene encoding for 4CL, isolated as described in example 1, was reamplified by PCR from the plasmid pESC-URA-4CL-VST (example 5) using the forward primer 5'-TG CCATGG CA

ATGGCGCCAC AAGAACAAGC AGTTT-3' SEQ ID NO 35 and the reverse

primer 5'-GC GGATCC CCT TCA CAA TCC ATT TGC TAG TTT TGCC-3' SEQ ID NO 36 with 5' overhangs containing the restriction

sites NcoI and BamHI, respectively. The introduction of restriction sites at the 5' and 3' ends of the gene allowed ligation of the restricted PCR product into a pET16b vector

(Novagen) digested with NcoI and BamHI. The resulting plasmid, pET16b-4CL, contained the gene encoding for 4CL under the control of the T7 promoter. Both the T7 promoter

and the gene encoding for 4CL were reamplified as one fragment by PCR from the plasmid pET16b-4CL using the

forward primer 5'-TT GCGGCCGC AAA TCT CGA TCC CGC GAA ATT AAT ACG-3' SEQ ID NO 37 and the reverse primer 5'-CG CTCGAG

CCT TCA CAA TCC ATT TGC TAG TTT TGCC-3' SEQ ID NO 38 with 5' overhangs, containing the restriction sites NotI and XhoI,

respectively. The introduction of restriction sites at the 5' and 3' ends of the DNA fragment allowed ligation of the

restricted PCR product into the plasmid pET26b-VST that was digested with NotI and XhoI before ligation. The resulting

plasmid, pET26b-VST-4CL, contained the two genes 4CL and VST that each were under control of an individual T7 promoter.

Example 18

Expression of the pathway to resveratrol in Escherichia coli, using TAL, 4CL and VST.

The transformation of the bacterial cell was conducted in accordance with methods known in the art, for instance, by using competent cells or by electroporation (see, e.g., Sambrook et al., 1989). The *E. coli* strain BL21 (DE3) (Novagen) was co-transformed with the two vectors pET16b-TAL (example 16) and pET26b-VST-4CL (Example 17), resulting in strain FSEC-TAL4CLVST. In addition, *E. coli* strain BL21 (DE3) was co-transformed with the two empty vectors pET16b (Novagen) and pET26b (Novagen), resulting in strain FSEC-control, which was used as a control strain. Transformants were selected on Luria-Bertani (LB) medium with 100 µg/ml ampicillin and 60 µg/ml kanamycin.

Example 19

Fermentation with recombinant Escherichia coli strains in shake flasks.

Pre-cultures of *Escherichia coli* BL21 (DE3) were grown in glass tubes at 160 rpm and 37 °C in 7 ml of LB medium containing 100 µg/ml ampicillin and 60 µg/ml kanamycin. Exponentially growing precultures were used for inoculation of 500 ml baffled shake flasks that contained 200 ml LB medium supplemented with 50 g/l glucose, 5 g/l K₂HPO₄, 80 µg/ml ampicillin and 50 µg/ml kanamycin, which were incubated at 160 rpm and 37 °C. After 5 hours, isopropyl β-thiogalactopyranoside (IPTG) was added at a final concentration of 1 mM, as an inducer of the T7 promoter that was in front of each of the three genes TAL, 4CL and VST. After an incubation period of 48 hours at 37 °C, the cells

were harvested and subjected to extraction procedures and analysed for the presence of produced resveratrol.

Example 20

5 *Extraction and analysis of resveratrol in Escherichia coli.*

Extraction and analysis was performed using the methods as described in example 14 and 15.

10 Results

Strain FSEC-TAL4CLVST and FSEC-control, were cultivated on 50 g/l glucose as described in example 19, and analyzed for their content of intracellular resveratrol according to
15 example 14 and 15. The HPLC-analysis showed that strain FSEC-TAL4CLVST did contain considerable amounts of a component with a retention time of 3.4 min., which is identical to coumaric acid (figure 6). However, the extract did not contain a component that eluted at the same time as
20 *trans*-resveratrol. Said result, therefore, indicated that the tyrosine ammonia lyase (TAL) was active indeed, but did not lead to production of detectable amounts of resveratrol. The lack of resveratrol formation, however, could be the result of; i) a non-functional coumarate-CoA ligase (4CL);
25 ii) a non-functional resveratrol synthase (VST); iii) too low levels of coumaric acid, caused by either non-optimal cultivation conditions, or non-optimal expression/activity of TAL, or branching of coumaric acid into other products. To evaluate said hypotheses, the strains were grown on
30 similar media as described in example 19 but now in the presence of 20 mg/l of coumaric acid. The subsequent HPLC-analysis of extracts of FSEC-TAL4CLVST indeed showed a cluster of peaks around the same retention time as *trans*-

resveratrol, which was not observed in extracts of FS-control (figure 6). Indeed, the UV absorption spectrum of the peak with a retention time of 5.5 min. was similar to the spectrum of pure *trans*-resveratrol (figure 7), whereas
5 no such spectrum could be obtained for peaks in the control strain. The results, therefore, strongly suggest the presence of an active phenylpropanoid pathway in *Escherichia coli*, which can lead to production of resveratrol. Most likely the production of resveratrol without addition of
10 coumaric acid can be achieved by cultivating the strains under well-defined growth conditions in batch- and continuous cultures, and/or optimizing the expression/activities of the individual enzymes.

15 Example 21

Construction of a bacterial vector for expression of PAL and C4H in Lactococcus lactis.

The plasmid pSH71 and derivatives thereof, which is used in
20 the following examples, is a bifunctional shuttle vector with multiple origins of replication from *Escherichia coli* and *Lactococcus lactis*. With that, the host range specificity traverses *Escherichia coli* and other species of lactic acid bacteria. Though transformations in *Lactococcus*
25 *lactis* usually proceed without problems, putative difficult transformations in other species of lactic acid bacteria can, therefore, be overcome by using *Escherichia coli* as an intermediate host for the construction of recombinant plasmids. The plasmid contains one or more marker genes to
30 allow the microorganism that harbour them to be selected from those which do not. The selection system that is used for *Lactococcus lactis* is based upon dominant markers, e.g. resistance against erythromycin and chloramphenicol, but

systems based upon genes involved in carbohydrate metabolism, peptidases and food grade markers, have also been described. In addition, the plasmid contains promoter- and terminator sequences that allow the expression of the recombinant genes. Suitable promoters are taken from genes of *Lactococcus lactis* e.g. *lacA*. Furthermore, the plasmid contains suitable unique restriction sites to facilitate the cloning of DNA fragments and subsequent identification of recombinants.

10 In the examples below the plasmid contains either the erythromycin resistance gene, designated as pSH71-ERY^r, or the chloramphenicol resistance gene, designated as pSH71-CM^r

The gene encoding PAL, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-PAL-C4H (example 3), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-ERY^r vector that contains the *lacA* promoter from *Lactococcus lactis*. The resulting plasmid, pSH71-ERY^r-PAL, contains the gene encoding PAL under the control of the *lacA* promoter from *Lactococcus lactis*.

The gene encoding C4H, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-PAL-C4H (example 3) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-CM^r vector to yield pSH71-CM^r-C4H. The *lacA* promoter and the gene encoding C4H are reamplified as one fragment by PCR from the plasmid pSH71-CM^r-C4H using forward- and reverse primers, with 5' overhangs containing

suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the DNA fragment allows ligation of the restricted PCR product into the digested plasmid pSH71-ERY^r-PAL. The resulting plasmid, pSH71-ERY^r-PAL-C4H, contains the genes encoding PAL and C4H that are each under the control of an individual *lacA* promoter. The sequence of the genes encoding PAL and C4H is verified by sequencing of two different clones of pSH71-ERY^r-PAL-C4H.

Example 22

Construction of a bacterial vector for expression of TAL in Lactococcus lactis.

The gene encoding for TAL, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-TAL (example 6) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-ERY^r vector. The resulting plasmid, pSH71-ERY^r-TAL, contains the gene encoding for TAL under the control of the *lacA* promoter from *Lactococcus lactis*. The sequence of the gene encoding for TAL is verified by sequencing of two different clones of pSH71-ERY^r-TAL.

Example 23

Construction of a bacterial vector for expression of 4CL and VST in Lactococcus lactis.

The gene encoding 4CL, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL-VST

(example 5), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-CM^r vector. The resulting plasmid, pSH71-CM^r-4CL, contains the gene encoding for 4CL under the control of the *lacA* promoter from *Lactobacillus lactis*.
The gene encoding VST, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL-VST (example 5) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-ERY^r vector. The resulting plasmid, pSH71-ERY^r-VST, contains the gene encoding VST under the control of the *lacA* promoter from *Lactococcus lactis*. The *lacA* promoter and the gene encoding VST are reamplified as one fragment by PCR from the plasmid pSH71-ERY^r-VST using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the DNA fragment allows ligation of the restricted PCR product into the digested plasmid pSH71-CM^r-4CL. The resulting plasmid, pSH71-CM^r-4CL-VST, contains the genes encoding 4CL and VST that are each under the control of their individual *lacA* promoter. The sequence of the genes encoding 4CL and VST is verified by sequencing of two different clones of pSH71-CM^r-4CL-VST.

Example 24

Expression of the pathway to resveratrol in *Lactococcus lactis*

- Lactococcus lactis* strains are transformed with the vectors described in examples 21, 22 and 23, separately or in combination. The transformation of the bacterial cell is conducted in accordance with methods known in the art, for instance, by using competent cells or by electroporation (see, e.g., Sambrook et al., 1989). Transformants are selected on medium containing the antibiotics erythromycin and chloramphenicol and streak purified on the same medium.
- 10 *Lactococcus lactis* strain MG1363 is transformed separately with the vector pSH71-ERY^r-TAL (example 22), yielding the strain FSLL-TAL; with pSH71-ERY^r-PAL-C4H (example 21), yielding the strain FSLL-PALC4H and with pSH71-CM^r-4CL-VST (example 23), yielding strain FSLL-4CLVST. In addition,
- 15 *Lactococcus lactis* strain MG1363 is co-transformed with pSH71-ERY^r-TAL (example 22) and pSH71-CM^r-4CL-VST (example 23), and the transformed strain is named FSLL-TAL4CLVST. The same strain is also co-transformed with pSH71-ERY^r-PAL-C4H (example 21), and pSH71-CM^r-4CL-VST (example 23), resulting
- 20 in the strain FSLL-PALC4H4CLVST.

Example 25

- 25 *Fermentation with recombinant Lactococcus lactis strains in fermentors.*

The recombinant yeast strains can be grown in fermenters operated as batch, fed-batch or chemostat cultures.

- 30 Batch and Fed-batch cultivations

The microorganism is grown in a baffled bioreactor with a working volume of 1.5 liters under anaerobic, aerobic or

microaerobic conditions. All cultures are incubated at 30°C, at 350 rpm. A constant pH of 6.6 is maintained by automatic addition of 10 M KOH. Cells are grown on lactose in defined MS10 medium supplemented with the following components to allow growth under aerobic conditions: MnSO₄ (1.25×10^{-5} g/l), thiamine (1 mg/l), and DL-6,8-thioctic acid (2.5 mg/l). The lactose concentration is, for example 50 g/l. The bioreactors are inoculated with cells from precultures grown at 30°C in shake flasks on the medium described above buffered with threefold-higher concentrations of K₂HPO₄ and KH₂PO₄. Anaerobic conditions are ensured by flushing the medium with N₂ (99.998% pure) prior to inoculation and by maintaining a constant flow of 50 ml/min of N₂ through the headspace of the bioreactor during cultivation. The bioreactors used for microaerobic and aerobic cultivation are equipped with polarographic oxygen sensors that are calibrated with air (DOT, 100%) and N₂ (DOT, 0%). Aerobic conditions are obtained by sparging the bioreactor with air at a rate of 1 vvm to ensure that the DOT is more than 80%. During microaerobic experiments the DOT is kept constant 5% by sparging the reactor with gas composed of a mixture of N₂ and atmospheric air, at a rate of 0.25 vvm.

Chemostat cultures

In chemostat cultures the cells can be grown in, for example, 1-L working-volume Applikon laboratory fermentors at 30 °C and 350 rpm. The dilution rate (*D*) can be set at different values, e.g. at 0.050 h⁻¹, 0.10 h⁻¹, 0.15 h⁻¹, or 0.20 h⁻¹. The pH is kept constant, e.g at 6.6, by automatic addition of 5 M KOH, using the growth medium described above, supplemented with antifoam (50 µl/l). The concentration of

lactose can be set at different values, e.g. is 3.0 g/l
6.0 g/l, 12.0 g/l, 15.0 g/l or 18.0 g/l. The bioreactor is
inoculated to an initial biomass concentration of 1 mg /l
and the feed pump is turned on at the end of the exponential
5 growth phase.

An anaerobic steady state is obtained by introducing
50 ml/min of N₂ (99.998% pure) into the headspace of the
bioreactor. Different anoxic steady states can obtained by
sparging the reactor with 250 ml/min of gas composed of N₂
10 (99.998% pure) and atmospheric air at various ratios. The
oxygen electrode is calibrated by sparging the bioreactor
with air (100% DOT) and with N₂ (0% DOT).

For all conditions, the gas is sterile filtered before being
introduced into the bioreactor. The off gas is led through a
15 condenser cooled to lower than -8°C and analyzed for its
volumetric content of CO₂ and O₂ by means of an acoustic gas
analyser.

Cultivations are considered to be in steady state after at
least 5 residence times, and if the concentrations of
20 biomass and fermentation end products remain unchanged (less
than 5% relative deviation) over the last two residence
times.

Example 26

25 *Extraction and analysis of resveratrol in Lactococcus lactis*

Extraction and analysis is performed using the methods as
described in examples 14 and 15.

30 Example 27

Construction of a fungal vector for expression of PAL and C4H in species belonging to the genus Aspergillus.

The plasmid that is used in the following examples, is
5 derived from pARp1 that contains the AMA1 initiating
replication sequence from *Aspergillus nidulans*, which also
sustains autonomous plasmid replication in *A. niger* and *A.*
oryzae (Gems et al., 1991). Moreover, the plasmid is a
shuttle vector, containing the replication sequence of
10 *Escherichia coli*, and the inherent difficult transformations
in *Aspergillus niger* and *Aspergillus oryzae* can therefore
overcome by using *Escherichia coli* as an intermediate host
for the construction of recombinant plasmids. The plasmid
contains one or more marker genes to allow the microorganism
15 that harbour them to be selected from those which do not.
The selection system can be either based upon dominant
markers e.g. resistance against hygromycin B, phleomycin and
bleomycin, or heterologous markers e.g amino acids and the
pyrG gene. In addition the plasmid contains promoter- and
20 terminator sequences that allow the expression of the
recombinant genes. Suitable promoters are taken from genes
of *Aspergillus nidulans* e.g. *alcA*, *glaA*, *amy*, *niaD*, and
gpdA. Furthermore, the plasmid contains suitable unique
restriction sites to facilitate the cloning of DNA fragments
25 and subsequent identification of recombinants.

The plasmid used in the following examples contains the
strong constitutive *gpdA*-promoter and auxotrophic markers,
all originating from *Aspergillus nidulans*; the plasmid
containing the gene *methG* that is involved in methionine
30 biosynthesis, is designated as pAMA1-MET; the plasmid
containing the gene *hisA* that is involved in histidine
biosynthesis, is designated as pAMA1-HIS.

The gene encoding PAL, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-PAL-C4H (example 3), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The
5 introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pAMA1-MET vector that contains the *gpdA* promoter from *Aspergillus nidulans*. The resulting plasmid, pAMA1-MET-PAL contains the gene encoding PAL under the
10 control of the *gpdA* promoter from *Aspergillus nidulans*.

The gene encoding C4H, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-PAL-C4H (example 3) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The
15 introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pAMA1-HIS vector to yield pAMA1-HIS-C4H. The *gpdA* promoter and the gene encoding C4H are reamplified as one fragment by PCR from the plasmid pAMA1-HIS-C4H using
20 forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the DNA fragment allows ligation of the restricted PCR product into the digested plasmid pAMA1-MET-PAL. The resulting plasmid,
25 pAMA1-MET-PAL-C4H, contains the genes encoding PAL and C4H that are each under the control of an individual *gpdA* promoter from *Aspergillus nidulans*. The sequence of the genes encoding PAL and C4H is verified by sequencing of two different clones of pAMA1-MET-PAL-C4H.

30

Example 28

Construction of a fungal vector for expression of TAL in species belonging to the genus Aspergillus.

The gene encoding for TAL, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-TAL (example 6) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pAMA1-MET vector. The resulting plasmid, pAMA1-MET-TAL, contains the gene encoding for TAL under the control of the *gpdA* promoter from *Aspergillus nidulans*. The sequence of the gene encoding for TAL is verified by sequencing of two different clones of pAMA1-MET-TAL.

15

Example 29

Construction of a fungal vector for expression of 4CL and VST in species belonging to the genus Aspergillus.

The gene encoding 4CL, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL-VST (example 5), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pAMA1-HIS vector that contains the *gpdA* promoter from *Aspergillus nidulans*. The resulting plasmid, pAMA1-HIS-4CL contains the gene encoding 4CL under the control of the *gpdA* promoter from *Aspergillus nidulans*.

The gene encoding VST, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL-VST (example 5) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The

introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pAMA1-MET vector to yield pAMA1-MET-VST. The *gpdA* promoter and the gene encoding VST are reamplified as
5 one fragment by PCR from the plasmid pAMA1-MET-VST using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the DNA fragment allows ligation of the restricted PCR product into the
10 digested plasmid pAMA1-HIS-4CL. The resulting plasmid, pAMA1-HIS-4CL-VST, contains the genes encoding 4CL and VST that are each under the control of an individual *gpdA* promoter from *Aspergillus nidulans*. The sequence of the genes encoding 4CL and VST is verified by sequencing of two
15 different clones of pAMA1-HIS-4CL-VST.

Example 30

Expression of the pathway to resveratrol in Aspergillus niger.

20 *Aspergillus niger* strains are transformed with the vectors described in examples 27, 28 and 29, separately or in combination. The transformation of the fungal cell is conducted in accordance with methods known in the art, for
25 instance, by electroporation or by conjugation (see, e.g., Sambrook et al., 1989). Transformants are selected on minimal medium lacking methionine and/or histidine.

A strain of *Aspergillus niger* that is auxotrophic for
30 histidine and methionine, for instance, strain FGSC A919 (see <http://www.fgsc.net>), is transformed separately with the vector pAMA1-MET-TAL (example 28), yielding the strain FSAN-TAL; with pAMA1-MET-PAL-C4H (example 27), yielding the

strain FSAN-PALC4H and with pAMA1-HIS-4CL-VST (example 29), yielding strain FSAN-4CLVST. In addition, *Aspergillus niger* strain FGSC A919 is co-transformed with pAMA1-MET-TAL (example 28) and pAMA1-HIS-4CL-VST (example 29), and the
5 transformed strain is named FSAN-TAL4CLVST. The same strain is also co-transformed with pAMA1-MET-PAL-C4H (example 27), and pAMA1-HIS-4CL-VST (example 29), resulting in the strain FSAN-PALC4H4CLVST.

10 Example 31

Expression of the pathway to resveratrol in Aspergillus oryzae.

A strain of *Aspergillus oryzae* that contains a native set of
15 genes encoding for PAL, C4H and 4CL (Seshime et al., 2005) and that is auxotrophic for methionine, is transformed with the vector pAMA1-MET-VST (example 29), yielding the strain FSAO-VST. The transformation of the fungal cell is conducted in accordance with methods known in the art, for instance,
20 by electroporation or by conjugation (see, e.g., Sambrook et al., 1989). Transformants are selected on minimal medium lacking methionine.

Example 32

25 *Fermentation with recombinant strains of Aspergillus niger and Aspergillus oryzae in fermentors.*

The recombinant yeast strains can be grown in fermenters operated as batch, fed-batch or chemostat cultures.

30

Batch and Fed-batch cultivations

The microorganism is grown in a baffled bioreactor with a working volume of 1.5 liters under aerobic conditions. All cultures are incubated at 30 °C, at 500 rpm. A constant pH of 6.0 is maintained by automatic addition of 10 M KOH, and aerobic conditions are obtained by sparging the bioreactor with air at a rate of 1 vvm to ensure that the DOT is more than 80%. Cells are grown on glucose in defined medium consisting of the following components to allow growth in batch cultivations: 7.3 g/l (NH₄)₂SO₄, 1.5 g/l KH₂PO₄, 1.0 g/l MgSO₄·7H₂O, 1.0 g/l NaCl, 0.1 g/l CaCl₂·2H₂O, 0.1 ml/l Sigma antifoam, 7.2 mg/l ZnSO₄·7H₂O, 1.3 mg/l CuSO₄·5H₂O, 0.3 mg/l NiCl₂·6H₂O, 3.5 mg/l MnCl₂·4H₂O and 6.9 mg/l FeSO₄·7H₂O. The glucose concentration is, for example, 10- 20-, 30-, 40- or 50 g/l. To allow growth in fed-batch cultivations the medium is composed of: 7.3 g/l (NH₄)₂SO₄, 4.0 g/l KH₂PO₄, 1.9 g/l MgSO₄·7H₂O, 1.3 g/l NaCl, 0.10 g/l CaCl₂·2H₂O, 0.1 ml/l Sigma antifoam, 7.2 mg/l ZnSO₄·7H₂O, 1.3 mg/l CuSO₄·5H₂O, 0.3 mg/l NiCl₂·6H₂O, 3.5 mg/l MnCl₂·4H₂O and 6.9 mg/l FeSO₄·H₂O in the batch phase. The reactor is then fed with, for example, 285 g/kg glucose and 42 g/kg (NH₄)₂SO₄.

Free mycelium from a pre-batch is used for inoculating the batch- and fed-batch cultures. A spore concentration of 2.10⁹ spores/l is used for inoculation of the pre-batch culture at pH 2.5. Spores are obtained by propagation of freeze-dried spores onto 29 g rice to which the following components are added: 6 ml 15 g/l sucrose, 2.3 g/l (NH₄)₂SO₄, 1.0 g/l KH₂PO₄, 0.5 g/l MgSO₄·7H₂O, 0.50 g/l NaCl, 14.3 mg/l ZnSO₄·7H₂O, 2.5 mg/ CuSO₄·5H₂O, 0.50 mg/l NiCl₂·6H₂O, and 13.8 mg/l FeSO₄·7H₂O. The spores are propagated at 30 °C for 7-14 days to yield a black layer of spores on the rice grains and are harvested by adding 100 ml of 0.1% Tween 20 in sterile water. For all conditions, the gas is sterile filtered before being introduced into the bioreactor. The off gas is

led through a condenser cooled to lower than -8°C and analyzed for its volumetric content of CO_2 and O_2 by means of an acoustic gas analyser.

5 Chemostat cultures

In chemostat cultures the cells can be grown in, for example, 1.5-L working-volume Biostat B laboratory fermentors at 30°C and 500 rpm. A constant pH of 6.0 is
10 maintained by automatic addition of 10 M KOH, and aerobic conditions are obtained by sparging the bioreactor with air at a rate of 1 vvm to ensure that the DOT is more than 80%. The dilution rate (D) can be set at different values, e.g. at 0.050 h^{-1} , 0.10 h^{-1} , 0.15 h^{-1} , or 0.20 h^{-1} . The pH is kept
15 constant, e.g. at 6.6, by automatic addition of 10 M KOH, using a minimal growth medium with the following components: 2.5 g/l $(\text{NH}_4)_2\text{SO}_4$, 0.75 g/l KH_2PO_4 , 1.0 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1.0 g/l NaCl, 0.1 g/l $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.1 ml/l Sigma antifoam, 7.2 mg/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 1.3 mg/l $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.3 mg/l $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, 3.5 mg/l
20 $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ and 6.9 mg/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$. The concentration of glucose can be set at different values, e.g. is 3.0 g/l, 6.0 g/l, 12.0 g/l, 15.0 g/l or 18.0 g/l. The bioreactor is inoculated with free mycelium from a pre-batch culture as described above, and the feed pump is turned on at the end
25 of the exponential growth phase.

For all conditions, the gas is sterile filtered before being introduced into the bioreactor. The off gas is led through a condenser cooled to lower than -8°C and analyzed for its volumetric content of CO_2 and O_2 by means of an acoustic gas
30 analyser.

Cultivations are considered to be in steady state after at least 5 residence times, and if the concentrations of

biomass glucose and composition of the off-gas remain unchanged (less than 5% relative deviation) over the last two residence times.

5 Example 33

Extraction and analysis of resveratrol in Aspergillus niger and Aspergillus oryzae

Extraction and analysis is performed using the methods as
10 described in examples 14 and 15.

REFERENCES

- Patent no. US-A-6521748
15 Patent no. US-A-2001053847
Patent no. US-A-2004059103
20 Patent no. US-A-2004023357

Allina, S.M., Pri-Hadash, A., Theilmann, D.A., Ellis, B.E. and Douglas, C.J. (1998) 4-coumarate: Coenzyme A ligase in hybrid poplar. Properties of enzymes, cDNA cloning, and
25 analysis of recombinant clones. Plant Physiol. 116, 743-754.

Aoyama, Y., Yoshida, Y., Kubota, S., Kumaoka, H. and Furumichi, A. (1978). NADPH-cytochrome P-450 reductase of yeast microsomes. Arch. Biochem. Biophys. 185, 362-369.

Becker JV, Armstrong GO, van der Merwe MJ, Lambrechts MG, Vivier MA, Pretorius IS. (2003). Metabolic engineering of *Saccharomyces cerevisiae* for the synthesis of the wine-related antioxidant resveratrol. FEMS Yeast Res. 4, 79-85.

Blanquet, S., Meunier, J. P., Minekus, M., Marol-Bonnin, S., and Alric, M. (2003). Recombinant *Saccharomyces cerevisiae* Expressing P450 in Artificial Digestive Systems: a Model for Biodegradation in the Human Digestive Environment. Appl. Environ. Microbiol. 69, 2884-2892.

Celotti E and others. (1996). Resveratrol content of some wines obtained from dried Valpolicella grapes: Recioto and Amarone. Journal of Chromatography A 730, 47-52.

Cochrane, F.C., Davin, L.B. and Lewis N.G. (2004). The *Arabidopsis* phenylalanine ammonia lyase gene family: kinetic characterization of the four PAL isoforms. Phytochemistry 65, 1557-1564.

Couzin, J. (2004) Aging Research's Family Feud. Science 303, 1276-1279.

Ehlting, J., Büttner, D., Wang, Q., Douglas, C.J., Somssich, I.E. and Kombrink, E. (1999). Three 4-coumarate:coenzyme A ligases in *Arabidopsis thaliana* represents two evolutionary divergent classes in angiosperms. The plant journal. 19, 9-20.

Filpula, D., Vaslet, C.A., Levy, A., Sykes, A. and Strausberg, R.L. Nucleotide sequence of gene for

phenylalanine ammonia-lyase from *Rhodotorula rubra*. (1988).
Nucleic Acids Res. 16, 11381.

5 Gehm, B.D., McAndrews, J.M., Chien, P.Y. and Jameson, J.L.
(1997). Resveratrol, a polyphenolic compound found in grapes
and wine, is an agonist for the estrogen receptor. Proc.
Natl. Acad. Sci. USA 94, 14138-14143.

10 Gems, D., Johnstone, I.L. and Clutterbuck, A.J. (1991). An
autonomously replicating plasmid transforms *Aspergillus*
nidulans at high frequency. Gene 98, 61-67.

Hain, R., Reif, H.J., Krause, E., Langebartels, R., Kindl,
H., Vornam, B., Wiese, W., Schmelzer, E., Schreier, P.H.,
Stocker, R.H. and Stenzel, K. (1993). Disease resistance
15 results from foreign phytoalexin expression in a novel
plant. Nature 361, 153-156.

Hwang EI, Kaneko M, Ohnishi Y, Horinouchi S. (2003).
Production of plant-specific flavanones by *Escherichia coli*
20 containing an artificial gene cluster. Appl. Environ.
Microbiol. 69, 2699-706.

Huang, M-T. (1997). Diet for cancer prevention. Food Sci.
24, 713-727

25 Hart, J. H. (1981) Annu. Rev. Phytopathology 19, 437-458.

Hart, J. H., Shrimpton, D. M. (1979) Phytopathology 69,
1138-1143.

30 Hall, S.S. (2003) In Vino Vitalis? Compounds Activate Life-
Extending Genes. Science 301, 1165.

Hamberger, B. and Hahlbrock, K. (2004). The 4-coumarate:CoA ligase gene family in *Arabidopsis thaliana* comprises one rare, sinapate-activating and three commonly occurring isoenzymes. Proc. Natl. Acad. Sci. USA. 101, 2209-2214.

5

Jang, M., Cai, L., Udeani, GO., Slowing, KV., Thomas, CF., Beecher, CWW., Fong, HHS., Farnsworth, NR., Kinghorn, AD., Mehta, RG., Moon, RC., Pezzuto, JM. (1997). Cancer Chemopreventive Activity of Resveratrol, a Natural Product
10 Derived from Grapes. Science 275, 218-220.

Jeandet, P. Bessis, R., Maume, B.F., Meunier, P., Peyron, D. and Trollat, P. (1995). Effect of Enological Practices on the Resveratrol Isomer Content of Wine J. Agric. Food Chem.
15 43, 316-319.

Jeandet, P. Bessis, R., Sbaghi, M. and Meunier, P. (1994). Occurrence of a resveratrol β -D-glucoside in wine: Preliminary studies. Vitis 33, 183-184.

20

Koopmann, E., Logemann, E. and Hahlbrock, K. (1999). Regulation and Functional Expression of Cinnamate 4-Hydroxylase from Parsley. Plant Physiol. 119, 49-55.

25 Kopp, P. (1998). Resveratrol, a phytoestrogen found in red wine. A possible explanation for the conundrum of the "French Paradox"? Eur. J. Endocrinol. 138, 619-620.

Kyndt JA, Meyer TE, Cusanovich MA, Van Beeumen JJ. (2002).
30 Characterization of a bacterial tyrosine ammonia lyase, a biosynthetic enzyme for the photoactive yellow protein. FEBS Lett. 512, 240-244.

- LaGrange, D.C., Pretorius, I.S. and Van Zyl, W.H. (1997). Cloning of the *Bacillus pumilus* beta-xylosidase gene (xynB) and its expression in *Saccharomyces cerevisiae*. Appl. Microbiol. Biotechnol. 47, 262-266.
- 5 Lin X., Kaul S., Rounsley S.D., Shea T.P., Benito M.-I., Town C.D., Fujii C.Y., Mason T.M., Bowman C.L., Barnstead M.E., Feldblyum T.V., Buell C.R., Ketchum K.A., Lee J.J., Ronning C.M., Koo H.L., Moffat K.S., Cronin L.A., Shen M., Pai G., Van Aken S., Umayam L., Tallon L.J., Gill J.E.,
- 10 Adams M.D., Carrera A.J., Creasy T.H., Goodman H.M., Somerville C.R., Copenhaver G.P., Preuss D., Nierman W.C., White O., Eisen J.A., Salzberg S.L., Fraser C.M., Venter J.C. (1999). Sequence and analysis of chromosome 2 of the plant *Arabidopsis thaliana*. Nature 402, 761-768.
- 15 Lobo, R.A. (1995). Benefits and risks of estrogen replacement therapy. Am. J. Obstet. Gynecol. 173, 982-989.
- Martin, V.J.J., Pitera, D.J., Withers, S.T., Newman, J.D.
- 20 and Keasling, J.D. (2003). Engineering a mevalonate pathway in *Escherichia coli* for production of terpenoids. Nature biotechnology 21, 796-802.
- Mizutani, M., Ohta, D. and Sato, R. (1997). Isolation of a
- 25 cDNA and a genomic clone encoding cinnamate 4-hydroxylase from *Arabidopsis* and its expression manner in planta. Plant Physiol. 113, 755-763.
- Ro, D.K., Mah, N., Ellis, B.E. and Douglas, C.J. (2001).
- 30 Functional Characterization and Subcellular Localization of Poplar (*Populus trichocarpa* x *Populus deltoides*) Cinnamate 4-Hydroxylase. Plant Physiol. 126, 317-329.

Ro D.K., Douglas C.J. (2004). Reconstitution of the entry point of plant phenylpropanoid metabolism in yeast (*Saccharomyces cerevisiae*): implications for control of metabolic flux into the phenylpropanoid pathway. J. Biol. Chem. 279, 2600-2607.

Rosler J, Krekel F, Amrhein N, Schmid J. (1997). Maize phenylalanine ammonia-lyase has tyrosine ammonia-lyase activity. Plant Physiol. 113, 175-179. activity. Plant physiol. 113, 175-179.

Samappito, S., Page, J.E., Schmidt, J., De-Eknamkul, W. and Kutchan, T.M. (2003). Aromatic and pyrone polyketides synthesized by a stilbene synthase from *Rheum tataricum*. Phytochemistry 62, 313-323.

Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). Molecular Cloning, A Laboratory Manual, 2d edition, Cold Spring Harbor, N.Y.

Schoppner, A.; Kindl, H. (1984) Purification and properties of a stilbene synthase from induced cell suspension cultures of peanut. J. Biol. Chem. 259, 6806-6811.

Seshime, Y., Juvvadi, P.R., Fujii, I. and Kitamoto, K. (2005). Genomic evidences for the existence of a phenylpropanoid metabolic pathway in *Aspergillus oryzae*. Biochem Biophys Res Commun. 337, 747-51.

Urban P., Mignotte, C., Kazmaier M., Delorme F. And Pompon D. (1997). Cloning, Yeast Expression, and Characterization

of the Coupling of Two Distantly Related *Arabidopsis thaliana* NADPH-Cytochrome 450 Reductases with P450 CYP73A5. J. Biol. Chem. 272, 19176-19186.

- 5 Watts, K.T., Lee, P.C. and Schmidt-Dannert, C. (2004). Exploring recombinant flavonoid biosynthesis in metabolically engineered *Escherichia coli*. Chembiochem 5, 500-507.

- 10 The following is a summary of the nucleotide and amino acid sequences appearing herein:

SEQ ID NO: 1 is a nucleotide sequence from *Arabidopsis thaliana* encoding a phenylalanine ammonia lyase (PAL2).

- 15 SEQ ID NO: 2 is the amino acid sequence encoded by SEQ ID NO: 1.

SEQ ID NO: 3 is a nucleotide sequence from *Arabidopsis thaliana* encoding a cinnamate 4-hydroxylase (C4H).

- 20 SEQ ID NO: 4 is the amino acid sequence encoded by SEQ ID NO: 3.

SEQ ID NO: 5 is a nucleotide sequence from *Arabidopsis thaliana* encoding a 4-coumarate:CoenzymeA ligase (4CL1).

SEQ ID NO: 6 is the amino acid sequence encoded by SEQ ID NO: 5.

- 25 SEQ ID NO: 7 is a nucleotide sequence from *Rheum tataricum* encoding a resveratrol synthase (VST).

SEQ ID NO: 8 is the amino acid sequence encoded by SEQ ID NO: 7.

- 30 SEQ ID NO: 9 is a nucleotide sequence from *Rheum tataricum* encoding a resveratrol synthase (VST), which is codon-optimized for expression in *S. cerevisiae*.

SEQ ID NO: 10 is the amino acid sequence encoded by SEQ ID NO: 9.

SEQ ID NO: 11 is a nucleotide sequence from *Rhodobacter capsulatus* encoding a tyrosine ammonia lyase (TAL).

SEQ ID NO: 12 is the amino acid sequence encoded by SEQ ID NO: 11.

- 5 SEQ ID NO: 13 is a nucleotide sequence from *Rhodobacter capsulatus* encoding a tyrosine ammonia lyase (TAL), which is codon-optimized for expression in *S. cerevisiae*.

SEQ ID NO: 14 is the amino acid sequence encoded by SEQ ID NO: 13.

- 10 SEQ ID NO: 15 is a nucleotide sequence from *S. cerevisiae* encoding a NADPH:cytochrome P450 reductase (CPR1).

SEQ ID NO: 16 is the amino acid sequence encoded by SEQ ID NO: 15.

- 15 SEQ ID NO: 17 is a nucleotide sequence from *Arabidopsis thalianus* encoding a NADPH:cytochrome P450 reductase (AR2).

SEQ ID NO: 18 is the amino acid sequence encoded by SEQ ID NO: 17.

SEQ ID NOs 19-32 are primer sequences appearing in Table 1, Example 1.

- 20 SEQ ID NOs 33-34 are primer sequences appearing in Example 16.

SEQ ID NOs 35-38 are primer sequences appearing in Example 17

SEQUENCE LISTING

<110> Fluxome Sciences A/S

5 <120> METABOLICALLY ENGINEERED CELLS FOR THE PRODUCTION OF RESVERATROL OR AN OLIGOMERIC OR GLYCOSIDICALLY-BOUND DERIVATIVE THEREOF

<130> PJS/P10686WO

10 <150> GB0503657.9
<151> 2005-02-22

<160> 38

15 <170> PatentIn version 3.3

<210> 1
<211> 2154
<212> DNA

20 <213> Arabidopsis thaliana

<400> 1

atggatcaaa tcgaagcaat gttgtgcggc ggaggagaga agacaaaagt ggcggttact	60
25 acgaagactt tggcagatcc attgaattgg ggttttagcag cggatcaaat gaaaggaagt	120
catttagatg aagtgaagaa gatggtcgaa gagtatcgta gaccagtcgt gaatcttggc	180
ggagaaacac tgacgatcgg acaagttgct gccatctcca ccgtaggagg cagcgttaag	240
30 gttgagttag cgggagacttc aagagccggt gtgaaagcta gcagtgattg ggttatggag	300
agcatgaaca aaggtactga cagttacgga gtcaccaccg gctttggtgc tacttctcac	360
35 cggagaacca aaaacggcac cgcattacaa acagaactca ttagattttt gaacgccgga	420
atattcggaa acacgaagga gacatgtcac aactgcccgc aatccgccac aagagccgcc	480
atgctcgtca gagtcaacac tcttctccaa ggatactccg ggatccgatt cgagatcctc	540
40 gaagcgatta caagtctcct caaccacaac atctctccgt cactacctct ccgtggaacc	600
attaccgcct ccggcgatct cgttctctctc tcttacatcg ccggacttct caccggccgt	660
45 cctaattcca aagccaccgg tcccgacggt gaatcgctaa ccgcgaaaga agcttttgag	720
aaagccggaa tcagtactgg attcttcgat ttacaacctt aggaagggtt agctctcggt	780
aatggcacgg cggttggatc tggaatggcg tcgatggttc tattcgaagc gaatgtccaa	840
50 gcggtgttag cggaggtttt atcagcgatc ttccgggagg ttatgagcgg gaaacctgag	900
tttaccgatc atctgactca tcgttttaaaa catcatcccg gacaaatcga agcggcgggc	960
55 ataatggagc acatactcga cggaagctca tacatgaaat tagctcaaaa ggttcacgag	1020
atggatccat tgcagaaacc aaaacaagat cgttacgctc ttctgtacatc tctcaatgg	1080
ctaggtcctc aaattgaagt aatccgtcaa gctacgaaat cgatagagcg tgaaatcaac	1140

	tccgttaacg ataatccggt gatcgatggt tcgaggaaca aggcgattca cggtaggtaac	1200
	ttccaaggaa caccaatcgg agtttctatg gataacacga gattggcgat tgctgcgatt	1260
5	gggaagctaa tgtttgctca atttctctgag cttgttaatg atttctacaa caatggactt	1320
	ccttcgaatc taactgcttc gagtaatcca agtttggatt atggattcaa aggagcagag	1380
10	attgctatgg cttcttattg ttctgagctt caatacttgg ctaatccagt cacaagccat	1440
	gttcaatcag ctgagcaaca taatcaagat gtgaactctc ttggtttgat ctctgtctgt	1500
	aaaacatctg aagctgtgga tattcttaag ctaatgtcaa caacgttcct tgtggggata	1560
15	tgtcaagctg ttgatttgag acatttggag gagaatctga gacaaactgt gaagaacaca	1620
	gtttctcaag ttgctaagaa agtgtaaac actggaatca acggtgagtt acatccgtca	1680
20	aggttttgcg agaaggactt gcttaagggt gttgatcgtg agcaagtgtt cacgtatgtg	1740
	gatgatcctt gtagcgctac gtaccggtg atgcagagac taagacaagt tattgttgat	1800
	cacgctttgt ccaacggtga gactgagaag aatgcagtga cttcgatctt tcaaaagatt	1860
25	ggagcttttg aagaggagct taaggctgtg cttccaaagg aagttgaagc ggctagagcg	1920
	gcttatggga atggaactgc gccgattcct aaccggatta aggaatgtag gtcgtatccg	1980
30	ttgtataggt tcgtgaggga agagcttgga acgaagttgt tgactggaga aaaggttggt	2040
	tctccgggag aggagtttga taaggtcttc actgctatgt gtgaaggtaa acttattgat	2100
35	ccgttgatgg attgtctcaa ggaatggaac ggagctccga ttccgatttg ctaa	2154
	<210> 2	
	<211> 717	
	<212> PRT	
40	<213> Arabidopsis thaliana	
	<400> 2	
45	Met Asp Gln Ile Glu Ala Met Leu Cys Gly Gly Gly Glu Lys Thr Lys	
	1 5 10 15	
	Val Ala Val Thr Thr Lys Thr Leu Ala Asp Pro Leu Asn Trp Gly Leu	
50	20 25 30	
	Ala Ala Asp Gln Met Lys Gly Ser His Leu Asp Glu Val Lys Lys Met	
	35 40 45	
55	Val Glu Glu Tyr Arg Arg Pro Val Val Asn Leu Gly Gly Glu Thr Leu	
	50 55 60	

60

	Thr	Ile	Gly	Gln	Val	Ala	Ala	Ile	Ser	Thr	Val	Gly	Gly	Ser	Val	Lys	65	70	75	80
5	Val	Glu	Leu	Ala	Glu	Thr	Ser	Arg	Ala	Gly	Val	Lys	Ala	Ser	Ser	Asp	85	90	95	
10	Trp	Val	Met	Glu	Ser	Met	Asn	Lys	Gly	Thr	Asp	Ser	Tyr	Gly	Val	Thr	100	105	110	
15	Thr	Gly	Phe	Gly	Ala	Thr	Ser	His	Arg	Arg	Thr	Lys	Asn	Gly	Thr	Ala	115	120	125	
	Leu	Gln	Thr	Glu	Leu	Ile	Arg	Phe	Leu	Asn	Ala	Gly	Ile	Phe	Gly	Asn	130	135	140	
20	Thr	Lys	Glu	Thr	Cys	His	Thr	Leu	Pro	Gln	Ser	Ala	Thr	Arg	Ala	Ala	145	150	155	160
25	Met	Leu	Val	Arg	Val	Asn	Thr	Leu	Leu	Gln	Gly	Tyr	Ser	Gly	Ile	Arg	165	170	175	
30	Phe	Glu	Ile	Leu	Glu	Ala	Ile	Thr	Ser	Leu	Leu	Asn	His	Asn	Ile	Ser	180	185	190	
35	Pro	Ser	Leu	Pro	Leu	Arg	Gly	Thr	Ile	Thr	Ala	Ser	Gly	Asp	Leu	Val	195	200	205	
40	Pro	Leu	Ser	Tyr	Ile	Ala	Gly	Leu	Leu	Thr	Gly	Arg	Pro	Asn	Ser	Lys	210	215	220	
	Ala	Thr	Gly	Pro	Asp	Gly	Glu	Ser	Leu	Thr	Ala	Lys	Glu	Ala	Phe	Glu	225	230	235	240
45	Lys	Ala	Gly	Ile	Ser	Thr	Gly	Phe	Phe	Asp	Leu	Gln	Pro	Lys	Glu	Gly	245	250	255	
50	Leu	Ala	Leu	Val	Asn	Gly	Thr	Ala	Val	Gly	Ser	Gly	Met	Ala	Ser	Met	260	265	270	
55	Val	Leu	Phe	Glu	Ala	Asn	Val	Gln	Ala	Val	Leu	Ala	Glu	Val	Leu	Ser	275	280	285	
	Ala	Ile	Phe	Ala	Glu	Val	Met	Ser	Gly	Lys	Pro	Glu	Phe	Thr	Asp	His	290	295	300	

Leu Thr His Arg Leu Lys His His Pro Gly Gln Ile Glu Ala Ala Ala
 305 310 315 320
 5
 Ile Met Glu His Ile Leu Asp Gly Ser Ser Tyr Met Lys Leu Ala Gln
 325 330 335
 10
 Lys Val His Glu Met Asp Pro Leu Gln Lys Pro Lys Gln Asp Arg Tyr
 340 345 350
 15
 Ala Leu Arg Thr Ser Pro Gln Trp Leu Gly Pro Gln Ile Glu Val Ile
 355 360 365
 20
 Arg Gln Ala Thr Lys Ser Ile Glu Arg Glu Ile Asn Ser Val Asn Asp
 370 375 380
 Asn Pro Leu Ile Asp Val Ser Arg Asn Lys Ala Ile His Gly Gly Asn
 385 390 395 400
 25
 Phe Gln Gly Thr Pro Ile Gly Val Ser Met Asp Asn Thr Arg Leu Ala
 405 410 415
 30
 Ile Ala Ala Ile Gly Lys Leu Met Phe Ala Gln Phe Ser Glu Leu Val
 420 425 430
 35
 Asn Asp Phe Tyr Asn Asn Gly Leu Pro Ser Asn Leu Thr Ala Ser Ser
 435 440 445
 40
 Asn Pro Ser Leu Asp Tyr Gly Phe Lys Gly Ala Glu Ile Ala Met Ala
 450 455 460
 Ser Tyr Cys Ser Glu Leu Gln Tyr Leu Ala Asn Pro Val Thr Ser His
 465 470 475 480
 45
 Val Gln Ser Ala Glu Gln His Asn Gln Asp Val Asn Ser Leu Gly Leu
 485 490 495
 50
 Ile Ser Ser Arg Lys Thr Ser Glu Ala Val Asp Ile Leu Lys Leu Met
 500 505 510
 55
 Ser Thr Thr Phe Leu Val Gly Ile Cys Gln Ala Val Asp Leu Arg His
 515 520 525
 Leu Glu Glu Asn Leu Arg Gln Thr Val Lys Asn Thr Val Ser Gln Val
 530 535 540

5	Ala Lys Lys Val Leu Thr Thr Gly Ile Asn Gly Glu Leu His Pro Ser 545 550 555 560
10	Arg Phe Cys Glu Lys Asp Leu Leu Lys Val Val Asp Arg Glu Gln Val 565 570 575
15	Phe Thr Tyr Val Asp Asp Pro Cys Ser Ala Thr Tyr Pro Leu Met Gln 580 585 590
20	Arg Leu Arg Gln Val Ile Val Asp His Ala Leu Ser Asn Gly Glu Thr 595 600 605
25	Glu Lys Asn Ala Val Thr Ser Ile Phe Gln Lys Ile Gly Ala Phe Glu 610 615 620
30	Glu Glu Leu Lys Ala Val Leu Pro Lys Glu Val Glu Ala Ala Arg Ala 625 630 635 640
35	Ala Tyr Gly Asn Gly Thr Ala Pro Ile Pro Asn Arg Ile Lys Glu Cys 645 650 655
40	Arg Ser Tyr Pro Leu Tyr Arg Phe Val Arg Glu Glu Leu Gly Thr Lys 660 665 670
45	Leu Leu Thr Gly Glu Lys Val Val Ser Pro Gly Glu Glu Phe Asp Lys 675 680 685
50	Val Phe Thr Ala Met Cys Glu Gly Lys Leu Ile Asp Pro Leu Met Asp 690 695 700
55	Cys Leu Lys Glu Trp Asn Gly Ala Pro Ile Pro Ile Cys 705 710 715
50	<210> 3 <211> 1518 <212> DNA <213> Arabidopsis thaliana
55	<400> 3 atggacctcc tcttgctgga gaagtcttta atcgccgtct tcgtggcggg gattctcgcc 60 acgggtgattt caaagctccg cggcaagaaa ttgaagctac ctccaggtcc tataccaatt 120 ccgatcttcg gaaactggct tcaagtcgga gatgatctca accaccgtaa tctcgtcgat 180 tacgctaaga aattcggcga tctcttcctc ctccgtatgg gtcagcgaaa cctagtcgtc 240

	gtctcctcac cggatctaac aaaggaagtg ctctcactc aaggcgttga gtttgatcc	300
	agaacgagaa acgtcgtgtt cgacattttc accgggaaaag gtcaagatat ggtgttcact	360
5	gtttacggcg agcattggag gaagatgaga agaatcatga cggttccttt cttaccaac	420
	aaagtgttc aacagaatcg tgaaggttg gagtttgaag cagctagtgt tgttgaagat	480
10	gttaagaaga atccagattc tgctacgaaa ggaatcgtgt tgaggaaaag tttgcaattg	540
	atgatgtata acaatatgtt ccgtatcatg ttcgatagaa gatttgagag tgaggatgat	600
	cctcttttcc ttaggcttaa ggctttgaat ggtgagagaa gtcgattagc tcagagcttt	660
15	gagtataact atggagattt cattcctatc cttagaccat tcctcagagg ctatttgaag	720
	atttgtcaag atgtgaaaga tcgaagaatc gctcttttca agaagtactt tgttgatgag	780
20	aggaagcaaa ttgcgagttc taagcctaca ggtagtgaag gattgaaatg tgccattgat	840
	cacatccttg aagctgagca gaaggagaa atcaacgagg acaatgttct ttacatcgtc	900
	gagaacatca atgtcgccgc gattgagaca acattgtggt ctatcgagtg gggaattgca	960
25	gagctagtga accatcctga aatccagagt aagctaagga acgaactcga cacagttctt	1020
	ggaccgggtg tgcaagtcac cgagcctgat cttcacaaac ttccatacct tcaagctgtg	1080
30	gttaaggaga ctcttcgtct gagaatggcg attcctctcc tcgtgcctca catgaacctc	1140
	catgatgcga agctcgctgg ctacgatatc ccagcagaaa gcaaaatcct tgttaatgct	1200
	tggtggctag caaacaaccc caacagctgg aagaagcctg aagagttag accagagagg	1260
35	ttctttgaag aagaatcgca cgtggaagct aacggtaatg acttcaggta tgtgccattt	1320
	ggtgttgac gtogaagctg tcccgggatt atattggcat tgcctatttt ggggatcacc	1380
40	attggtagga tggccagaa cttcgagctt cttcctctc caggacagtc taaagtggat	1440
	actagtgaga aaggtggaca attcagcttg cacatcctta accactccat aatcgttatg	1500
45	aaaccaagga actgttaa	1518
	<210> 4	
	<211> 505	
	<212> PRT	
50	<213> Arabidopsis thaliana	
	<400> 4	
55	Met Asp Leu Leu Leu Leu Glu Lys Ser Leu Ile Ala Val Phe Val Ala	
	1 5 10 15	
	Val Ile Leu Ala Thr Val Ile Ser Lys Leu Arg Gly Lys Lys Leu Lys	
	20 25 30	

	Leu	Pro	Pro	Gly	Pro	Ile	Pro	Ile	Phe	Gly	Asn	Trp	Leu	Gln		
5			35					40			45					
	Val	Gly	Asp	Asp	Leu	Asn	His	Arg	Asn	Leu	Val	Asp	Tyr	Ala	Lys	Lys
		50					55					60				
10																
	Phe	Gly	Asp	Leu	Phe	Leu	Leu	Arg	Met	Gly	Gln	Arg	Asn	Leu	Val	Val
	65					70					75					80
15																
	Val	Ser	Ser	Pro	Asp	Leu	Thr	Lys	Glu	Val	Leu	Leu	Thr	Gln	Gly	Val
					85					90					95	
20																
	Glu	Phe	Gly	Ser	Arg	Thr	Arg	Asn	Val	Val	Phe	Asp	Ile	Phe	Thr	Gly
				100					105					110		
25																
	Lys	Gly	Gln	Asp	Met	Val	Phe	Thr	Val	Tyr	Gly	Glu	His	Trp	Arg	Lys
			115					120					125			
30																
	Met	Arg	Arg	Ile	Met	Thr	Val	Pro	Phe	Phe	Thr	Asn	Lys	Val	Val	Gln
		130					135					140				
35																
	Gln	Asn	Arg	Glu	Gly	Trp	Glu	Phe	Glu	Ala	Ala	Ser	Val	Val	Glu	Asp
	145					150					155					160
40																
	Val	Lys	Lys	Asn	Pro	Asp	Ser	Ala	Thr	Lys	Gly	Ile	Val	Leu	Arg	Lys
					165					170					175	
45																
	Arg	Leu	Gln	Leu	Met	Met	Tyr	Asn	Asn	Met	Phe	Arg	Ile	Met	Phe	Asp
				180					185					190		
50																
	Arg	Arg	Phe	Glu	Ser	Glu	Asp	Asp	Pro	Leu	Phe	Leu	Arg	Leu	Lys	Ala
			195					200					205			
55																
	Leu	Asn	Gly	Glu	Arg	Ser	Arg	Leu	Ala	Gln	Ser	Phe	Glu	Tyr	Asn	Tyr
		210					215					220				
60																
	Gly	Asp	Phe	Ile	Pro	Ile	Leu	Arg	Pro	Phe	Leu	Arg	Gly	Tyr	Leu	Lys
	225					230					235					240
65																
	Ile	Cys	Gln	Asp	Val	Lys	Asp	Arg	Arg	Ile	Ala	Leu	Phe	Lys	Lys	Tyr
					245					250					255	
70																
	Phe	Val	Asp	Glu	Arg	Lys	Gln	Ile	Ala	Ser	Ser	Lys	Pro	Thr	Gly	Ser

65

					260					265					270				
5	Glu	Gly	Leu	Lys	Cys	Ala	Ile	Asp	His	Ile	Leu	Glu	Ala	Glu	Gln	Lys			
			275					280					285						
10	Gly	Glu	Ile	Asn	Glu	Asp	Asn	Val	Leu	Tyr	Ile	Val	Glu	Asn	Ile	Asn			
		290					295					300							
15	Val	Ala	Ala	Ile	Glu	Thr	Thr	Leu	Trp	Ser	Ile	Glu	Trp	Gly	Ile	Ala			
	305					310					315					320			
20	Glu	Leu	Val	Asn	His	Pro	Glu	Ile	Gln	Ser	Lys	Leu	Arg	Asn	Glu	Leu			
				325						330					335				
25	Asp	Thr	Val	Leu	Gly	Pro	Gly	Val	Gln	Val	Thr	Glu	Pro	Asp	Leu	His			
			340						345					350					
30	Lys	Leu	Pro	Tyr	Leu	Gln	Ala	Val	Val	Lys	Glu	Thr	Leu	Arg	Leu	Arg			
			355					360					365						
35	Met	Ala	Ile	Pro	Leu	Leu	Val	Pro	His	Met	Asn	Leu	His	Asp	Ala	Lys			
		370					375					380							
40	Leu	Ala	Gly	Tyr	Asp	Ile	Pro	Ala	Glu	Ser	Lys	Ile	Leu	Val	Asn	Ala			
	385					390					395					400			
45	Trp	Trp	Leu	Ala	Asn	Asn	Pro	Asn	Ser	Trp	Lys	Lys	Pro	Glu	Glu	Phe			
				405						410					415				
50	Arg	Pro	Glu	Arg	Phe	Phe	Glu	Glu	Glu	Ser	His	Val	Glu	Ala	Asn	Gly			
				420					425					430					
55	Asn	Asp	Phe	Arg	Tyr	Val	Pro	Phe	Gly	Val	Gly	Arg	Arg	Ser	Cys	Pro			
		435					440					445							
60	Gly	Ile	Ile	Leu	Ala	Leu	Pro	Ile	Leu	Gly	Ile	Thr	Ile	Gly	Arg	Met			
		450					455					460							
65	Val	Gln	Asn	Phe	Glu	Leu	Leu	Pro	Pro	Pro	Gly	Gln	Ser	Lys	Val	Asp			
	465					470					475					480			
70	Thr	Ser	Glu	Lys	Gly	Gly	Gln	Phe	Ser	Leu	His	Ile	Leu	Asn	His	Ser			
				485						490					495				

Ile Ile Val Met Lys Pro Arg Asn Cys
500 505

5	<210> 5		
	<211> 1686		
	<212> DNA		
	<213> Arabidopsis thaliana		
10	<400> 5		
	atggcgccac aagaacaagc agttttctcag gtgatggaga aacagagcaa caacaacaac	60	
	agtgacgtca ttttccgatc aaagttaccg gatatttaca tcccgaacca cctatctctc	120	
15	caogactaca ttttccaaaa catctccgaa ttcgccacta agccttgccct aatcaacgga	180	
	ccaaccggcc acgtgtacac ttactccgac gtccacgtca tctcccgcca aatcgccgcc	240	
	aattttcaca aactcggcgt taacccaaac gacgtcgtca tgctcctcct cccaaactgt	300	
20	ccogaattcg tctctctctt cctcgccgcc tcttccgcg gcgcaaccgc caccgccgca	360	
	aaccctttct tcaactccggc ggagatagct aaacaagcca aagcctccaa caccaaaactc	420	
25	ataatcaccg aagctcgtta cgtcgacaaa atcaaaccac ttcaaaacga cgacggagta	480	
	gtcatcgtct gcacgcagca caacgaatcc gtgccaatcc ctgaaggctg cctccgcttc	540	
	accgagttga ctacgtcgac aaccgaggca tcagaagtca tcgactcggg ggagatttca	600	
30	cgggacgacg tgggtggcact accttactcc tctggcacga cgggattacc aaaaggagtg	660	
	atgctgactc acaagggact agtcacgagc gttgctcagc aagtcgacgg cgagaaccgc	720	
35	aatctttatt tocacagcga tgacgtcata ctctgtgttt tgcccatggt tcatatctac	780	
	gctttgaact cgatcatggt gtgtggtctt agagttggtg cggcgattct gataatgcg	840	
	aagtttgaga tcaatctgct attggagctg atccagaggt gtaaagtac ggtggctccg	900	
40	atggttccgc cgattgtgtt ggccattgcy aagtcttcgg agacggagaa gtatgatttg	960	
	agctcgataa gagtgggtgaa atctggtgct gctcctcttg gtaaagaact tgaagatgcc	1020	
45	gttaatgcc aagtttctaa tgccaaactc ggtcagggat acggaatgac ggaagcaggt	1080	
	ccagtgcctag caatgtcgtt aggttttgca aaggaacctt ttccggttaa gtcaggagct	1140	
	tgtggtactg ttgtaagaaa tgctgagatg aaaatagttg atccagacac cggagattct	1200	
50	ctttcgagga atcaaccgcg tgagatttgt attcgtggtc accagatcat gaaaggttac	1260	
	ctcaacaatc cggcagctac agcagagacc attgataaag acggttggtc tcatactgga	1320	
55	gatattggat tgatcgatga cgatgacgag cttttcatcg ttgatcgatt gaaagaactt	1380	
	atcaagtata aaggttttca ggtagctccg gctgagctag aggctttgct catcggtcat	1440	
	cctgacatta ctgatgttgc tgttgcgcga atgaaagaag aagcagctgg tgaagttcct	1500	

gttgcatttg tggtgaaatc gaaggattcg gagttatcag aagatgatgt gaagcaattc 1560
 gtgtcgaaac aggttgtgtt ttacaagaga atcaacaaag tgttcttcac tgaatccatt 1620
 5 cctaaagctc catcaggga gatattgagg aaagatctga gggcaaaact agcaaatgga 1680
 ttgtga 1686
 10
 <210> 6
 <211> 561
 <212> PRT
 <213> Arabidopsis thaliana
 15
 <400> 6
 Met Ala Pro Gln Glu Gln Ala Val Ser Gln Val Met Glu Lys Gln Ser
 1 5 10 15
 20 Asn Asn Asn Asn Ser Asp Val Ile Phe Arg Ser Lys Leu Pro Asp Ile
 20 25 30
 25 Tyr Ile Pro Asn His Leu Ser Leu His Asp Tyr Ile Phe Gln Asn Ile
 35 40 45
 30 Ser Glu Phe Ala Thr Lys Pro Cys Leu Ile Asn Gly Pro Thr Gly His
 50 55 60
 35 Val Tyr Thr Tyr Ser Asp Val His Val Ile Ser Arg Gln Ile Ala Ala
 65 70 75 80
 40 Asn Phe His Lys Leu Gly Val Asn Gln Asn Asp Val Val Met Leu Leu
 85 90 95
 45 Leu Pro Asn Cys Pro Glu Phe Val Leu Ser Phe Leu Ala Ala Ser Phe
 100 105 110
 50 Arg Gly Ala Thr Ala Thr Ala Ala Asn Pro Phe Phe Thr Pro Ala Glu
 115 120 125
 55 Ile Ala Lys Gln Ala Lys Ala Ser Asn Thr Lys Leu Ile Ile Thr Glu
 130 135 140
 Ala Arg Tyr Val Asp Lys Ile Lys Pro Leu Gln Asn Asp Asp Gly Val
 145 150 155 160
 Val Ile Val Cys Ile Asp Asp Asn Glu Ser Val Pro Ile Pro Glu Gly
 165 170 175

5 Cys Leu Arg Phe Thr Glu Leu Thr Gln Ser Thr Thr Glu Ala Ser Glu
 180 185 190
 Val Ile Asp Ser Val Glu Ile Ser Pro Asp Asp Val Val Ala Leu Pro
 195 200 205
 10 Tyr Ser Ser Gly Thr Thr Gly Leu Pro Lys Gly Val Met Leu Thr His
 210 215 220
 15 Lys Gly Leu Val Thr Ser Val Ala Gln Gln Val Asp Gly Glu Asn Pro
 225 230 235 240
 20 Asn Leu Tyr Phe His Ser Asp Asp Val Ile Leu Cys Val Leu Pro Met
 245 250 255
 25 Phe His Ile Tyr Ala Leu Asn Ser Ile Met Leu Cys Gly Leu Arg Val
 260 265 270
 Gly Ala Ala Ile Leu Ile Met Pro Lys Phe Glu Ile Asn Leu Leu Leu
 275 280 285
 30 Glu Leu Ile Gln Arg Cys Lys Val Thr Val Ala Pro Met Val Pro Pro
 290 295 300
 35 Ile Val Leu Ala Ile Ala Lys Ser Ser Glu Thr Glu Lys Tyr Asp Leu
 305 310 315 320
 40 Ser Ser Ile Arg Val Val Lys Ser Gly Ala Ala Pro Leu Gly Lys Glu
 325 330 335
 45 Leu Glu Asp Ala Val Asn Ala Lys Phe Pro Asn Ala Lys Leu Gly Gln
 340 345 350
 Gly Tyr Gly Met Thr Glu Ala Gly Pro Val Leu Ala Met Ser Leu Gly
 355 360 365
 50 Phe Ala Lys Glu Pro Phe Pro Val Lys Ser Gly Ala Cys Gly Thr Val
 370 375 380
 55 Val Arg Asn Ala Glu Met Lys Ile Val Asp Pro Asp Thr Gly Asp Ser
 385 390 395 400
 Leu Ser Arg Asn Gln Pro Gly Glu Ile Cys Ile Arg Gly His Gln Ile

69

	405	410	415
5	Met Lys Gly Tyr Leu Asn Asn Pro Ala Ala Thr Ala Glu Thr Ile Asp 420 425 430		
10	Lys Asp Gly Trp Leu His Thr Gly Asp Ile Gly Leu Ile Asp Asp Asp 435 440 445		
15	Asp Glu Leu Phe Ile Val Asp Arg Leu Lys Glu Leu Ile Lys Tyr Lys 450 455 460		
20	Gly Phe Gln Val Ala Pro Ala Glu Leu Glu Ala Leu Leu Ile Gly His 465 470 475 480		
25	Pro Asp Ile Thr Asp Val Ala Val Val Ala Met Lys Glu Glu Ala Ala 485 490 495		
30	Gly Glu Val Pro Val Ala Phe Val Val Lys Ser Lys Asp Ser Glu Leu 500 505 510		
35	Ser Glu Asp Asp Val Lys Gln Phe Val Ser Lys Gln Val Val Phe Tyr 515 520 525		
40	Lys Arg Ile Asn Lys Val Phe Phe Thr Glu Ser Ile Pro Lys Ala Pro 530 535 540		
45	Ser Gly Lys Ile Leu Arg Lys Asp Leu Arg Ala Lys Leu Ala Asn Gly 545 550 555 560		
50	Leu		
55	<210> 7 <211> 1176 <212> DNA <213> Rheum tataricum		
60	<400> 7 atggcaccgg aggagtccag gcatgctgaa actgcagtta acagagccgc caccgtcctg		60
120	gccatcggca ctgccaaacc gccaaactgc tactatcaag cggactttcc tgactttctac		120
180	ttccgtgcc acaacagcga ccacctcacg cacctcaagc aaaaatttaa gcgcatttgt		180
240	gagaaatcga tgattgaaaa acgttatctc catttgacgg aagaaattot caaggagaat		240
300	ccaaatattg cttccttcga ggcgccatca ttggatgtaa gacataacat tcaagtgaaa		300

gaagtgggtgc tgctcggaag agaggcagct ttgaaggcca tcaatgagtg gggccaaccc 360
 aagtcaaaga tcacgcgcct catttgtgtg tgtattgccg gcgttgacat gcccggcgca 420
 5 gactatcaac tcaactaaact ccttggtcta caactttctg ttaagcgatt tatgttttac 480
 cacctaggat gctatgccg tggcaccgtc cttcgccctg cgaaggacat agcagaaaac 540
 aacaaggaag ctcggtgttct catcgttcgc tctgagatga cgccaatctg tttccgtggg 600
 10 ccatccgaaa ccacataga ctccatggtg gggcaagcaa tatttggtga cgggtgctgcg 660
 gctgttatag ttggtgcaaa tcccgacctg tccatcgaaa ggccgatttt cgagttgatt 720
 15 tctacatccc aaactatcat acctgaatcc gatgggtgcga ttgagggaca tttgcttgaa 780
 gttggactca gtttccaact ctaccagact gttccctcat taatctctaa ttgtatcgaa 840
 acttgtcttt caaaggcttt cacacctctt aacattagtg attggaactc actattctgg 900
 20 attgcacacc ctgggtggccg tgctatcctt gacgatatcg aggctactgt tgggtctcaag 960
 aaggagaaac ttaaggcaac aagacaagtt ttgaacgact atgggaacat gtcaagtgtc 1020
 25 tgcgatatttt tcatcatgga tgagatgagg aagaagtcgc tcgcaaacgg tcaagtaacc 1080
 actggagaag gactcaagtg ggggtgttctt tttgggttcg ggccaggtgt tactgtggaa 1140
 actgtggttc taagcagtgt gccgctaatt acctga 1176
 30

<210> 8

<211> 391

<212> PRT

35 <213> Rheum tataricum

<400> 8

40 Met Ala Pro Glu Glu Ser Arg His Ala Glu Thr Ala Val Asn Arg Ala
 1 5 10 15

45 Ala Thr Val Leu Ala Ile Gly Thr Ala Asn Pro Pro Asn Cys Tyr Tyr
 20 25 30

Gln Ala Asp Phe Pro Asp Phe Tyr Phe Arg Ala Thr Asn Ser Asp His
 35 40 45

50 Leu Thr His Leu Lys Gln Lys Phe Lys Arg Ile Cys Glu Lys Ser Met
 50 55 60

55 Ile Glu Lys Arg Tyr Leu His Leu Thr Glu Glu Ile Leu Lys Glu Asn
 65 70 75 80

Pro Asn Ile Ala Ser Phe Glu Ala Pro Ser Leu Asp Val Arg His Asn

71

		85		90		95
5	Ile Gln Val Lys Glu Val Val Leu Leu Gly Lys Glu Ala Ala Leu Lys	100		105		110
10	Ala Ile Asn Glu Trp Gly Gln Pro Lys Ser Lys Ile Thr Arg Leu Ile	115		120		125
15	Val Cys Cys Ile Ala Gly Val Asp Met Pro Gly Ala Asp Tyr Gln Leu	130		135		140
20	Thr Lys Leu Leu Gly Leu Gln Leu Ser Val Lys Arg Phe Met Phe Tyr	145		150		155
25	His Leu Gly Cys Tyr Ala Gly Gly Thr Val Leu Arg Leu Ala Lys Asp	165		170		175
30	Ile Ala Glu Asn Asn Lys Glu Ala Arg Val Leu Ile Val Arg Ser Glu	180		185		190
35	Met Thr Pro Ile Cys Phe Arg Gly Pro Ser Glu Thr His Ile Asp Ser	195		200		205
40	Met Val Gly Gln Ala Ile Phe Gly Asp Gly Ala Ala Ala Val Ile Val	210		215		220
45	Gly Ala Asn Pro Asp Leu Ser Ile Glu Arg Pro Ile Phe Glu Leu Ile	225		230		235
50	Ser Thr Ser Gln Thr Ile Ile Pro Glu Ser Asp Gly Ala Ile Glu Gly	245		250		255
55	His Leu Leu Glu Val Gly Leu Ser Phe Gln Leu Tyr Gln Thr Val Pro	260		265		270
	Ser Leu Ile Ser Asn Cys Ile Glu Thr Cys Leu Ser Lys Ala Phe Thr	275		280		285
	Pro Leu Asn Ile Ser Asp Trp Asn Ser Leu Phe Trp Ile Ala His Pro	290		295		300
	Gly Gly Arg Ala Ile Leu Asp Asp Ile Glu Ala Thr Val Gly Leu Lys	305		310		315
						320

	Lys	Glu	Lys	Leu	Lys	Ala	Thr	Arg	Gln	Val	Leu	Asn	Asp	Tyr	Gly	Asn	
					325					330					335		
5	Met	Ser	Ser	Ala	Cys	Val	Phe	Phe	Ile	Met	Asp	Glu	Met	Arg	Lys	Lys	
				340					345					350			
10	Ser	Leu	Ala	Asn	Gly	Gln	Val	Thr	Thr	Gly	Glu	Gly	Leu	Lys	Trp	Gly	
			355					360					365				
15	Val	Leu	Phe	Gly	Phe	Gly	Pro	Gly	Val	Thr	Val	Glu	Thr	Val	Val	Leu	
		370					375					380					
20	Ser	Ser	Val	Pro	Leu	Ile	Thr										
	385					390											
25	<210>	9															
	<211>	1176															
	<212>	DNA															
	<213>	Rheum tataricum															
30	<400>	9															
	atg	gccccag	aag	agagcag	gcacgcagaa	acggccg	tta	acagagctgc	aactgtttt	g							60
	gct	attggta	cggccaatcc	acccaattgt	tactatcaag	ctgactttcc	tgatttttat										120
	ttc	agagcca	caa	atagcga	tcatttgact	catcttaagc	aaaaatttaa	aaggatatgc									180
	gag	aagtcca	tgattgaaaa	gagatacttg	caccttaccg	aagagatctt	aaaagaaaac										240
35	cca	aatatag	cttcttttga	agctccctcc	ttagatgtac	gtcacaacat	tcaagtcaag										300
	gag	gtggttt	tacttggtaa	ggaagccgca	ttgaaagcta	taaacgaatg	gggacagcct										360
	aaa	agtaaga	taaccagatt	gatcgtatgt	tgc	atagctg	gcgttgacat	gcctgggtgca									420
40	gatt	atcaac	taacaaaatt	gctgggtcta	caattatccg	taaaaagg	tt	tatgttctac									480
	catt	taggct	gttacgctgg	tggcacagtt	ttaagactgg	ctaaggatat	agcagaaaat										540
45	aaca	aggagg	ctagagtctt	aatagtgcgt	agtgaaatga	ctcctatttg	ctttagaggt										600
	ccat	cagaaa	cacatatcga	cagcatggta	ggtcaggcaa	ttttcgg	tga	tggtgctgca									660
	gcc	gtaattg	tgggagctaa	tcctgattta	agtatcgaaa	gacctatttt	tgaacttatt										720
50	tct	acttcgc	aaaccattat	ccccgaatca	gatgggtgcaa	ttgaaggcca	tttattggag										780
	gtt	ggtttgt	cctttcaatt	gtatcagaca	gtgccatctt	taatttcaaa	ctgtatagaa										840
55	acct	gtctaa	gtaaagcatt	tacaccatta	aacattttctg	actggaattc	tttgttctgg										900
	att	gctcatc	caggtggaag	agccatctta	gatgacatcg	aagctactgt	gggactgaaa										960
	aag	gaaaaaac	taaaagctac	tagacaagtt	ttaa	atgact	acg	gtaatat	gtcatctgct								1020

tgtgtatattt tcattatgga tgagatgaga aaaaagtcac ttgcaaattgg ccaggtcacg 1080
 acaggtgagg gtctaaaatg gggagtccta ttcggttcg gccaggtgt cactgttgaa 1140
 5 accgttgtcc tgtcttcggt tccattgatc acttaa 1176

<210> 10
 10 <211> 391
 <212> PRT
 <213> Rheum tataricum

<400> 10
 15 Met Ala Pro Glu Glu Ser Arg His Ala Glu Thr Ala Val Asn Arg Ala
 1 5 10 15

20 Ala Thr Val Leu Ala Ile Gly Thr Ala Asn Pro Pro Asn Cys Tyr Tyr
 20 25 30

25 Gln Ala Asp Phe Pro Asp Phe Tyr Phe Arg Ala Thr Asn Ser Asp His
 35 40 45

30 Leu Thr His Leu Lys Gln Lys Phe Lys Arg Ile Cys Glu Lys Ser Met
 50 55 60

35 Ile Glu Lys Arg Tyr Leu His Leu Thr Glu Glu Ile Leu Lys Glu Asn
 65 70 75 80

40 Pro Asn Ile Ala Ser Phe Glu Ala Pro Ser Leu Asp Val Arg His Asn
 85 90 95

45 Ile Gln Val Lys Glu Val Val Leu Leu Gly Lys Glu Ala Ala Leu Lys
 100 105 110

50 Ala Ile Asn Glu Trp Gly Gln Pro Lys Ser Lys Ile Thr Arg Leu Ile
 115 120 125

55 Val Cys Cys Ile Ala Gly Val Asp Met Pro Gly Ala Asp Tyr Gln Leu
 130 135 140

Thr Lys Leu Leu Gly Leu Gln Leu Ser Val Lys Arg Phe Met Phe Tyr
 145 150 155 160

His Leu Gly Cys Tyr Ala Gly Gly Thr Val Leu Arg Leu Ala Lys Asp
 165 170 175

74

	Ile	Ala	Glu	Asn	Asn	Lys	Glu	Ala	Arg	Val	Leu	Ile	Val	Arg	Ser	Glu	
				180					185					190			
5	Met	Thr	Pro	Ile	Cys	Phe	Arg	Gly	Pro	Ser	Glu	Thr	His	Ile	Asp	Ser	
			195					200					205				
10	Met	Val	Gly	Gln	Ala	Ile	Phe	Gly	Asp	Gly	Ala	Ala	Ala	Val	Ile	Val	
		210					215					220					
15	Gly	Ala	Asn	Pro	Asp	Leu	Ser	Ile	Glu	Arg	Pro	Ile	Phe	Glu	Leu	Ile	
	225					230					235					240	
20	Ser	Thr	Ser	Gln	Thr	Ile	Ile	Pro	Glu	Ser	Asp	Gly	Ala	Ile	Glu	Gly	
					245					250					255		
25	His	Leu	Leu	Glu	Val	Gly	Leu	Ser	Phe	Gln	Leu	Tyr	Gln	Thr	Val	Pro	
				260					265					270			
30	Ser	Leu	Ile	Ser	Asn	Cys	Ile	Glu	Thr	Cys	Leu	Ser	Lys	Ala	Phe	Thr	
			275					280					285				
35	Pro	Leu	Asn	Ile	Ser	Asp	Trp	Asn	Ser	Leu	Phe	Trp	Ile	Ala	His	Pro	
		290					295					300					
40	Gly	Gly	Arg	Ala	Ile	Leu	Asp	Asp	Ile	Glu	Ala	Thr	Val	Gly	Leu	Lys	
	305					310					315					320	
45	Lys	Glu	Lys	Leu	Lys	Ala	Thr	Arg	Gln	Val	Leu	Asn	Asp	Tyr	Gly	Asn	
					325					330					335		
50	Met	Ser	Ser	Ala	Cys	Val	Phe	Phe	Ile	Met	Asp	Glu	Met	Arg	Lys	Lys	
				340					345					350			
55	Ser	Leu	Ala	Asn	Gly	Gln	Val	Thr	Thr	Gly	Glu	Gly	Leu	Lys	Trp	Gly	
			355					360					365				
60	Val	Leu	Phe	Gly	Phe	Gly	Pro	Gly	Val	Thr	Val	Glu	Thr	Val	Val	Leu	
		370					375					380					
65	Ser	Ser	Val	Pro	Leu	Ile	Thr										
	385					390											

<210> 11
 <211> 1596
 <212> DNA

<213> Rhodobacter capsulatus

<400> 11

5	atgaccctgc agtcacagac ggccaaggac tgcctcgcgc tggacggggc gctgacactt	60
	gtccaatgcg aggccatcgc gacacatcgc agccggattt cggtgacccc cgcgctgcgc	120
	gagcgtgcgc cgcgggcccc tgcgggctt gagcacgcca tcgccgagca gcgccacatt	180
10	tacggcatca ccaccggctt cggcccgtg gcgaaccgtc tgatcggggc cgatcagggg	240
	gcggagctgc agcagaacct gatctatcat ctggccaccg gcgtcggggc gaaactgagc	300
15	tgggcccagg cgcgggctt gatgctggcg cggctcaact cgatcctgca aggcgcgtcg	360
	ggggcctcgc cggagacgat cgaccggatc gttgcggtgc tcaatgcggg gtttgcccc	420
	gaggttcctg cgcagggaac ggtggggcgc tcgggcgatc tgaccccgtc tgcgcatatg	480
20	gtgctggcgc tgcagggacg ggggcggatg atcgaccctc cgggcccgtc gcaggaggcc	540
	ggggcggtga tggatcggct ctgcggcggt ccgctgacgc tggcgggccg tgacgggctg	600
	gcgctggtga atggcacctc ggcgatgacc gcgattgcgg ccctgaccgg ggtcgaggcg	660
25	gcgcgggcga tcgacgcgcg gcttcggcac agcgcggtcc tgatggaggc cttgtccggt	720
	catgccgaag cctggcatcc ggctttcgca gagctgcgcc cgcacccggg gcagctgcgg	780
30	gcgaccgagc ggctggcgca ggcgctggat ggggcggggc gggctctgtc gacctgacc	840
	gcggcgcggc ggctgaccgc cgcggatctg cggcccgaag atcatccggc gcaggatgcc	900
	tacagtctgc gcgtgggtgc gcaactggtc ggcgcggtct gggacacgct ggactggcac	960
35	gatcgtgtcg tcacctgcga gctcaattcc gtcaccgaca atccgatctt tcccagggc	1020
	tgcgcggtgc ccgccctgca cggcggcaat ttcattggcg tgcatgtcgc cttgacctc	1080
40	gatgcgtga acgcggcgct ggtgacgctg gcgggccttg tcgagcgtca gatcggcccg	1140
	ctgaccgacg aaaagctgaa caagggcctg cccgccttcc tgcacggggg gcaggcgggg	1200
45	ctgcaatcgg gcttcattgg ggcgaggtc acggcgacgg cgcttctggc ggaaatgcgg	1260
	gcgaatgcca cgcgggttc ggtgcagtc ctgtcgacca atggcgccaa tcaggatgtg	1320
	gtctcgatgg gaacgattgc cgcgcggagg gcgcgggcgc agctgtgcc cctgtcgag	1380
50	atccaggcga tcctggcgct tgcccttgcc caggcgatgg atctgcttga cgaccccgag	1440
	gggcaggccg gatggtcgtc tacggcgcgg gatctgcggg accggatccg ggcggtctcg	1500
55	cccgggcttc gcgccgacag accgcttgcc gggcatatcg aagcggtggc acagggtctg	1560
	cgtcacccct ccgcgcgcgc cgatcccccg gcatga	1596

<210> 12

<211> 531
 <212> PRT
 <213> Rhodobacter capsulatus

5 <400> 12

Met Thr Leu Gln Ser Gln Thr Ala Lys Asp Cys Leu Ala Leu Asp Gly
 1 5 10 15

10

Ala Leu Thr Leu Val Gln Cys Glu Ala Ile Ala Thr His Arg Ser Arg
 20 25 30

15

Ile Ser Val Thr Pro Ala Leu Arg Glu Arg Cys Ala Arg Ala His Ala
 35 40 45

20

Arg Leu Glu His Ala Ile Ala Glu Gln Arg His Ile Tyr Gly Ile Thr
 50 55 60

25

Thr Gly Phe Gly Pro Leu Ala Asn Arg Leu Ile Gly Ala Asp Gln Gly
 65 70 75 80

30

Ala Glu Leu Gln Gln Asn Leu Ile Tyr His Leu Ala Thr Gly Val Gly
 85 90 95

Pro Lys Leu Ser Trp Ala Glu Ala Arg Ala Leu Met Leu Ala Arg Leu
 100 105 110

35

Asn Ser Ile Leu Gln Gly Ala Ser Gly Ala Ser Pro Glu Thr Ile Asp
 115 120 125

40

Arg Ile Val Ala Val Leu Asn Ala Gly Phe Ala Pro Glu Val Pro Ala
 130 135 140

45

Gln Gly Thr Val Gly Ala Ser Gly Asp Leu Thr Pro Leu Ala His Met
 145 150 155 160

Val Leu Ala Leu Gln Gly Arg Gly Arg Met Ile Asp Pro Ser Gly Arg
 165 170 175

50

Val Gln Glu Ala Gly Ala Val Met Asp Arg Leu Cys Gly Gly Pro Leu
 180 185 190

55

Thr Leu Ala Ala Arg Asp Gly Leu Ala Leu Val Asn Gly Thr Ser Ala
 195 200 205

Met Thr Ala Ile Ala Ala Leu Thr Gly Val Glu Ala Ala Arg Ala Ile

	210	215	220	
5	Asp Ala Ala Leu Arg His Ser Ala Val Leu Met Glu Val Leu Ser Gly 225 230 235 240			
10	His Ala Glu Ala Trp His Pro Ala Phe Ala Glu Leu Arg Pro His Pro 245 250 255			
15	Gly Gln Leu Arg Ala Thr Glu Arg Leu Ala Gln Ala Leu Asp Gly Ala 260 265 270			
20	Gly Arg Val Cys Arg Thr Leu Thr Ala Ala Arg Arg Leu Thr Ala Ala 275 280 285			
25	Asp Leu Arg Pro Glu Asp His Pro Ala Gln Asp Ala Tyr Ser Leu Arg 290 295 300			
30	Val Val Pro Gln Leu Val Gly Ala Val Trp Asp Thr Leu Asp Trp His 305 310 315 320			
35	Asp Arg Val Val Thr Cys Glu Leu Asn Ser Val Thr Asp Asn Pro Ile 325 330 335			
40	Phe Pro Glu Gly Cys Ala Val Pro Ala Leu His Gly Gly Asn Phe Met 340 345 350			
45	Gly Val His Val Ala Leu Ala Ser Asp Ala Leu Asn Ala Ala Leu Val 355 360 365			
50	Thr Leu Ala Gly Leu Val Glu Arg Gln Ile Ala Arg Leu Thr Asp Glu 370 375 380			
55	Lys Leu Asn Lys Gly Leu Pro Ala Phe Leu His Gly Gly Gln Ala Gly 385 390 395 400			
	Leu Gln Ser Gly Phe Met Gly Ala Gln Val Thr Ala Thr Ala Leu Leu 405 410 415			
	Ala Glu Met Arg Ala Asn Ala Thr Pro Val Ser Val Gln Ser Leu Ser 420 425 430			
	Thr Asn Gly Ala Asn Gln Asp Val Val Ser Met Gly Thr Ile Ala Ala 435 440 445			

	Arg Arg Ala Arg Ala Gln Leu Leu Pro Leu Ser Gln Ile Gln Ala Ile	
	450 455 460	
5	Leu Ala Leu Ala Leu Ala Gln Ala Met Asp Leu Leu Asp Asp Pro Glu	
	465 470 475 480	
10	Gly Gln Ala Gly Trp Ser Leu Thr Ala Arg Asp Leu Arg Asp Arg Ile	
	485 490 495	
15	Arg Ala Val Ser Pro Gly Leu Arg Ala Asp Arg Pro Leu Ala Gly His	
	500 505 510	
20	Ile Glu Ala Val Ala Gln Gly Leu Arg His Pro Ser Ala Ala Ala Asp	
	515 520 525	
25	Pro Pro Ala	
	530	
30	<210> 13 <211> 1596 <212> DNA <213> Rhodobacter capsulatus	
35	<400> 13 atgaccctgc aatctcaaac agctaaagat tgtttggtt tggatggtgc cttgacatta	60
	gttcaatgcg aagcgatagc aacccataga agtagaatct ctgtaacacc agccctacgt	120
40	gagagatgtg ctagagcaca tgctagggtta gaacatgcaa tagccgaaca gcgacacata	180
	tatgggataa cgacaggctt cgggccactt gctaacaggc tgatcggagc agaccagggt	240
45	gctgaattac aacagaacct tatctaccat ttggcaaccg gagttggccc caaattatca	300
	tgggccgaag ccagagcttt aatgctcgct cgtttgaata gtatactaca aggtgcttct	360
50	ggtgctagcc ctgaaacaat tgataggatc gttgcagtct taaatgccgg atttgccccg	420
	gaagtcccg ccgaaggaac cgttggtgct tcgggtgact taactccggt agcacacatg	480
55	gtattagcat tgcaaggcag aggtcgtatg attgatcctt caggagaggt tcaagaagcc	540
	ggcgctgtca tggatagggt gtgtggaggc cctttaacat tggctgccag agatggcctc	600
	gccttagtaa atggtacatc tgccatgaca gctattgccg cattgaccgg tgtggaggct	660
	gcaagagcga ttgatgcagc gcttagacat tccgcagtct tgatggagggt cctgtcaggg	720
	catgctgagg cttggcaccg tgcctttgcg gaattgcgtc cgcacccagg acaattacgc	780
	gccactgaga ggtagctca agcattggac ggcgaggta gagtctgccg gactcttaca	840
	gccgctaggc gtctaactgc agctgatctg agaccagaag atcatccagc tcaagatgca	900

5 tattcacttc gagtagttcc tcagctgggt ggtgccgtat gggatacgtt ggattggcac 960
 gacaggggtt tgacttgcca acttaactcc gtgaccgaca atccaatttt ccccgagggt 1020
 tgtgcgggtc cagcactaca cggtggaac tttatgggcg tacatgtggc actagcttct 1080
 gacgctttaa atgcagcgtt gggtacatta gctggcttag ttgaaaggca gattgcaaga 1140
 10 cttactgatg agaagttgaa taagggtttg cctgcttttt tgcattggagg ccaagcagggt 1200
 ttacaatcag gtttcattgg agctcagggt actgctactg ctttgctagc ggaaatgaga 1260
 gctaacgcga ctcccggtgc cgttcaaagc ctgagcacca atgggtgcaa tcaagacgtg 1320
 15 gtaagtatgg gtacgattgc cgcgagacga gcaagagctc aacttttacc tctgtctcaa 1380
 atccaagcga ttttggcact ggctcttgca caagccatgg atctcctaga cgatcctgaa 1440
 20 ggacaagccg gttggtcctt aacggcaaga gatttaagag accgtatacg ggctgtcagt 1500
 ccaggggttc gcgcagatag accactagcg ggtcatattg aagctgtggc tcaaggtcta 1560
 agacaccctt cggcagctgc cgatccacct gcttaa 1596
 25

<210> 14
 <211> 531
 <212> PRT
 30 <213> Rhodobacter capsulatus

<400> 14

35 Met Thr Leu Gln Ser Gln Thr Ala Lys Asp Cys Leu Ala Leu Asp Gly
 1 5 10 15

40 Ala Leu Thr Leu Val Gln Cys Glu Ala Ile Ala Thr His Arg Ser Arg
 20 25 30

45 Ile Ser Val Thr Pro Ala Leu Arg Glu Arg Cys Ala Arg Ala His Ala
 35 40 45

Arg Leu Glu His Ala Ile Ala Glu Gln Arg His Ile Tyr Gly Ile Thr
 50 55 60

50 Thr Gly Phe Gly Pro Leu Ala Asn Arg Leu Ile Gly Ala Asp Gln Gly
 65 70 75 80

55 Ala Glu Leu Gln Gln Asn Leu Ile Tyr His Leu Ala Thr Gly Val Gly
 85 90 95

Pro Lys Leu Ser Trp Ala Glu Ala Arg Ala Leu Met Leu Ala Arg Leu
 100 105 110

5 Asn Ser Ile Leu Gln Gly Ala Ser Gly Ala Ser Pro Glu Thr Ile Asp
 115 120 125

10 Arg Ile Val Ala Val Leu Asn Ala Gly Phe Ala Pro Glu Val Pro Ala
 130 135 140

15 Gln Gly Thr Val Gly Ala Ser Gly Asp Leu Thr Pro Leu Ala His Met
 145 150 155 160

20 Val Leu Ala Leu Gln Gly Arg Gly Arg Met Ile Asp Pro Ser Gly Arg
 165 170 175

25 Val Gln Glu Ala Gly Ala Val Met Asp Arg Leu Cys Gly Gly Pro Leu
 180 185 190

30 Thr Leu Ala Ala Arg Asp Gly Leu Ala Leu Val Asn Gly Thr Ser Ala
 195 200 205

35 Met Thr Ala Ile Ala Ala Leu Thr Gly Val Glu Ala Ala Arg Ala Ile
 210 215 220

40 Asp Ala Ala Leu Arg His Ser Ala Val Leu Met Glu Val Leu Ser Gly
 225 230 235 240

45 His Ala Glu Ala Trp His Pro Ala Phe Ala Glu Leu Arg Pro His Pro
 245 250 255

50 Gly Gln Leu Arg Ala Thr Glu Arg Leu Ala Gln Ala Leu Asp Gly Ala
 260 265 270

55 Gly Arg Val Cys Arg Thr Leu Thr Ala Ala Arg Arg Leu Thr Ala Ala
 275 280 285

60 Asp Leu Arg Pro Glu Asp His Pro Ala Gln Asp Ala Tyr Ser Leu Arg
 290 295 300

65 Val Val Pro Gln Leu Val Gly Ala Val Trp Asp Thr Leu Asp Trp His
 305 310 315 320

70 Asp Arg Val Val Thr Cys Glu Leu Asn Ser Val Thr Asp Asn Pro Ile
 325 330 335

75 Phe Pro Glu Gly Cys Ala Val Pro Ala Leu His Gly Gly Asn Phe Met

81

		340		345		350	
5	Gly Val His Val Ala Leu Ala Ser Asp Ala Leu Asn Ala Ala Leu Val	355		360		365	
10	Thr Leu Ala Gly Leu Val Glu Arg Gln Ile Ala Arg Leu Thr Asp Glu	370		375		380	
15	Lys Leu Asn Lys Gly Leu Pro Ala Phe Leu His Gly Gly Gln Ala Gly	385		390		395	400
20	Leu Gln Ser Gly Phe Met Gly Ala Gln Val Thr Ala Thr Ala Leu Leu		405		410		415
25	Ala Glu Met Arg Ala Asn Ala Thr Pro Val Ser Val Gln Ser Leu Ser		420		425		430
30	Thr Asn Gly Ala Asn Gln Asp Val Val Ser Met Gly Thr Ile Ala Ala		435		440		445
35	Arg Arg Ala Arg Ala Gln Leu Leu Pro Leu Ser Gln Ile Gln Ala Ile		450		455		460
40	Leu Ala Leu Ala Leu Ala Gln Ala Met Asp Leu Leu Asp Asp Pro Glu		465		470		475
45	Gly Gln Ala Gly Trp Ser Leu Thr Ala Arg Asp Leu Arg Asp Arg Ile		485		490		495
50	Arg Ala Val Ser Pro Gly Leu Arg Ala Asp Arg Pro Leu Ala Gly His		500		505		510
55	Ile Glu Ala Val Ala Gln Gly Leu Arg His Pro Ser Ala Ala Ala Asp		515		520		525
	Pro Pro Ala	530					
	<210> 15						
	<211> 2076						
	<212> DNA						
	<213> Saccharomyces cerevisiae						
	<400> 15						
	atgccgtttg gaatagacaa caccgacttc actgtcctgg cggggctagt gcttgccgtg						

	ctactgtacg taaagagaaa ctccatcaag gaactgctga tgtccgatga cggagatata	120
	acagctgtca gctcgggcaa cagagacatt gctcaggtgg tgaccgaaaa caacaagaac	180
5	tacttggtgt tgtatgcgc gcagactggg actgccgagg attacgcca aaagttttcc	240
	aaggagctgg tggccaagtt caacctaaac gtgatgtgcg cagatgttga gaactacgac	300
10	tttgagtcgc taaacgatgt gcccgtcata gtctcgattt ttatctctac atatggtgaa	360
	ggagacttcc ccgacggggc ggtcaacttt gaagacttta tttgtaatgc ggaagcgggt	420
	gcactatcga acctgaggta taatatgttt ggtctgggaa attctactta tgaattcttt	480
15	aatggtgccg ccaagaaggc cgagaagcat ctctccgcg cgggcgctat cagactaggc	540
	aagctcgggtg aagctgatga tgggtgcagga actacagacg aagattacat ggctggaag	600
20	gactccatcc tggaggtttt gaaagacgaa ctgcatttgg acgaacagga agccaagttc	660
	acctctcaat tccagtacac tgtgttgaac gaaatcactg actccatgtc gcttggtgaa	720
	ccctctgctc actatttgcc ctgcacacg ttgaaccgca acgcagacgg catccaattg	780
25	ggtcccttcg atttgtctca accgtatatt gcacccatcg tgaaatctcg cgaactgttc	840
	tcttccaatg accgtaattg catccactct gaatttgact tgtccggctc taacatcaag	900
30	tactccactg gtgaccatct tgctgtttgg ccttccaacc cattggaaaa ggtcgaacag	960
	ttcttatcca tattcaacct ggacctgaa accatttttg acttgaagcc cctggatccc	1020
	accgtcaaag tgcccttccc aacgccaact actattggcg ctgctattaa acactatttg	1080
35	gaaattacag gacctgtctc cagacaattg ttttcatctt tgattcagtt cgcctccaac	1140
	gctgacgtca aggaaaaatt gactctgctt tcgaaagaca aggaccaatt cgcctcgag	1200
40	ataacctcca aatatttcaa catcgcagat gctctgaaat atttgtctga tggcgccaaa	1260
	tgggacaccg taccatgca attcttggtc gaatcagttc cccaaatgac tcctcgttac	1320
	tactctatct ctctctctc totgtctgaa aagcaaaccg tccatgtcac ctccattgtg	1380
45	gaaaactttc ctaaccaga attgcctgat gctcctccag ttgttggtgt tacgactaac	1440
	ttgttaagaa acattcaatt ggctcaaac aatgttaaca ttgccgaaac taacctacct	1500
50	gttcactacg atttaaattg ccacgtaaa ctttctgcca attacaaatt gcccgccac	1560
	gttcgtcgtt ctaacttcag attgccttcc aaccttcca cccagttat catgatcgg	1620
	ccaggtagcg gtgttgcccc attccgtggg tttatcagag agcgtgtcgc gttcctcgaa	1680
55	tcacaaaaga agggcggtaa caacgtttcg ctaggtaagc atatactgtt ttatggatcc	1740
	cgtaacactg atgatttctt gtaccaggac gaatggccag aatacgccaa aaaattggat	1800
	ggttcgttcg aaatggctcg ggccattcc aggttgccaa acaccaaaaa agtttatgtt	1860

caagataaat taaaggatta cgaagaccaa gtatttgaaa tgattaacaa cgggtgcattt 1920
 atctacgtct gtggtgatgc aaaggggatg gccaaagggtg tgtcaaccgc attggttggc 1980
 5 atcttatccc gtggtaaatc cattaccact gatgaagcaa cagagctaata caagatgctc 2040
 aagacttcag gtagatacca agaagatgctc tggtaa 2076
 10
 <210> 16
 <211> 691
 <212> PRT
 <213> *Saccharomyces cerevisiae*
 15
 <400> 16
 Met Pro Phe Gly Ile Asp Asn Thr Asp Phe Thr Val Leu Ala Gly Leu
 1 5 10 15
 20 Val Leu Ala Val Leu Leu Tyr Val Lys Arg Asn Ser Ile Lys Glu Leu
 20 25 30
 25 Leu Met Ser Asp Asp Gly Asp Ile Thr Ala Val Ser Ser Gly Asn Arg
 35 40 45
 30 Asp Ile Ala Gln Val Val Thr Glu Asn Asn Lys Asn Tyr Leu Val Leu
 50 55 60
 35 Tyr Ala Ser Gln Thr Gly Thr Ala Glu Asp Tyr Ala Lys Lys Phe Ser
 65 70 75 80
 Lys Glu Leu Val Ala Lys Phe Asn Leu Asn Val Met Cys Ala Asp Val
 85 90 95
 40 Glu Asn Tyr Asp Phe Glu Ser Leu Asn Asp Val Pro Val Ile Val Ser
 100 105 110
 45 Ile Phe Ile Ser Thr Tyr Gly Glu Gly Asp Phe Pro Asp Gly Ala Val
 115 120 125
 50 Asn Phe Glu Asp Phe Ile Cys Asn Ala Glu Ala Gly Ala Leu Ser Asn
 130 135 140
 55 Leu Arg Tyr Asn Met Phe Gly Leu Gly Asn Ser Thr Tyr Glu Phe Phe
 145 150 155 160
 Asn Gly Ala Ala Lys Lys Ala Glu Lys His Leu Ser Ala Ala Gly Ala
 165 170 175

5 Ile Arg Leu Gly Lys Leu Gly Glu Ala Asp Asp Gly Ala Gly Thr Thr
 180 185 190

10 Asp Glu Asp Tyr Met Ala Trp Lys Asp Ser Ile Leu Glu Val Leu Lys
 195 200 205

15 Asp Glu Leu His Leu Asp Glu Gln Glu Ala Lys Phe Thr Ser Gln Phe
 210 215 220

20 Gln Tyr Thr Val Leu Asn Glu Ile Thr Asp Ser Met Ser Leu Gly Glu
 225 230 235 240

25 Pro Ser Ala His Tyr Leu Pro Ser His Gln Leu Asn Arg Asn Ala Asp
 245 250 255

30 Gly Ile Gln Leu Gly Pro Phe Asp Leu Ser Gln Pro Tyr Ile Ala Pro
 260 265 270

35 Ile Val Lys Ser Arg Glu Leu Phe Ser Ser Asn Asp Arg Asn Cys Ile
 275 280 285

40 His Ser Glu Phe Asp Leu Ser Gly Ser Asn Ile Lys Tyr Ser Thr Gly
 290 295 300

45 Asp His Leu Ala Val Trp Pro Ser Asn Pro Leu Glu Lys Val Glu Gln
 305 310 315 320

50 Phe Leu Ser Ile Phe Asn Leu Asp Pro Glu Thr Ile Phe Asp Leu Lys
 325 330 335

55 Pro Leu Asp Pro Thr Val Lys Val Pro Phe Pro Thr Pro Thr Thr Ile
 340 345 350

60 Gly Ala Ala Ile Lys His Tyr Leu Glu Ile Thr Gly Pro Val Ser Arg
 355 360 365

65 Gln Leu Phe Ser Ser Leu Ile Gln Phe Ala Pro Asn Ala Asp Val Lys
 370 375 380

70 Glu Lys Leu Thr Leu Leu Ser Lys Asp Lys Asp Gln Phe Ala Val Glu
 385 390 395 400

75 Ile Thr Ser Lys Tyr Phe Asn Ile Ala Asp Ala Leu Lys Tyr Leu Ser

85

				405					410					415					
5	Asp	Gly	Ala	Lys	Trp	Asp	Thr	Val	Pro	Met	Gln	Phe	Leu	Val	Glu	Ser			
				420					425					430					
10	Val	Pro	Gln	Met	Thr	Pro	Arg	Tyr	Tyr	Ser	Ile	Ser	Ser	Ser	Ser	Leu			
			435					440					445						
15	Ser	Glu	Lys	Gln	Thr	Val	His	Val	Thr	Ser	Ile	Val	Glu	Asn	Phe	Pro			
			450				455					460							
20	Asn	Pro	Glu	Leu	Pro	Asp	Ala	Pro	Pro	Val	Val	Gly	Val	Thr	Thr	Asn			
	465					470					475					480			
25	Leu	Leu	Arg	Asn	Ile	Gln	Leu	Ala	Gln	Asn	Asn	Val	Asn	Ile	Ala	Glu			
				485						490					495				
30	Thr	Asn	Leu	Pro	Val	His	Tyr	Asp	Leu	Asn	Gly	Pro	Arg	Lys	Leu	Phe			
			500						505					510					
35	Ala	Asn	Tyr	Lys	Leu	Pro	Val	His	Val	Arg	Arg	Ser	Asn	Phe	Arg	Leu			
			515					520					525						
40	Pro	Ser	Asn	Pro	Ser	Thr	Pro	Val	Ile	Met	Ile	Gly	Pro	Gly	Thr	Gly			
		530					535					540							
45	Val	Ala	Pro	Phe	Arg	Gly	Phe	Ile	Arg	Glu	Arg	Val	Ala	Phe	Leu	Glu			
	545					550					555					560			
50	Ser	Gln	Lys	Lys	Gly	Gly	Asn	Asn	Val	Ser	Leu	Gly	Lys	His	Ile	Leu			
				565					570						575				
55	Phe	Tyr	Gly	Ser	Arg	Asn	Thr	Asp	Asp	Phe	Leu	Tyr	Gln	Asp	Glu	Trp			
			580					585						590					
60	Pro	Glu	Tyr	Ala	Lys	Lys	Leu	Asp	Gly	Ser	Phe	Glu	Met	Val	Val	Ala			
			595					600					605						
65	His	Ser	Arg	Leu	Pro	Asn	Thr	Lys	Lys	Val	Tyr	Val	Gln	Asp	Lys	Leu			
		610					615						620						
70	Lys	Asp	Tyr	Glu	Asp	Gln	Val	Phe	Glu	Met	Ile	Asn	Asn	Gly	Ala	Phe			
	625					630					635					640			

	Ile Tyr Val Cys Gly Asp Ala Lys Gly Met Ala Lys Gly Val Ser Thr	
	645	650 655
5	Ala Leu Val Gly Ile Leu Ser Arg Gly Lys Ser Ile Thr Thr Asp Glu	
	660	665 670
10	Ala Thr Glu Leu Ile Lys Met Leu Lys Thr Ser Gly Arg Tyr Gln Glu	
	675	680 685
15	Asp Val Trp	
	690	
	<210> 17	
	<211> 2136	
	<212> DNA	
20	<213> Arabidopsis thaliana	
	<400> 17	
	atgtcctctt cttcttcttc gtcaacctcc atgatcgatc tcatggcagc aatcatcaaa	60
25	ggagagcctg taattgtctc cgaccagct aatgcctccg cttacgagtc cgtagctgct	120
	gaattatcct ctatgcttat agagaatcgt caattcgcca tgattgttac cacttccatt	180
	gctgttctta ttggttgcat cgttatgctc gtttgaggga gatccggttc tgggaattca	240
30	aaacgtgtcg agcctcttaa gcctttgggtt attaagcctc gtgaggaaga gattgatgat	300
	gggcgtaaga aagttacat ctttttcggt acacaaactg gtactgctga aggttttgca	360
35	aaggcttttag gagaagaagc taaagcaaga tatgaaaaga ccagattcaa aatcgttgat	420
	ttggatgatt acgcggctga tgatgatgag tatgaggaga aattgaagaa agaggatgtg	480
	gctttcttct tcttagccac atatggagat ggtgagccta ccgacaatgc agcgagattc	540
40	tacaaatggt tcaccgaggg gaatgacaga ggagaatggc ttaagaactt gaagtatgga	600
	gtgtttggat taggaaacag acaatatgag cattttaata aggttgccaa agttgtagat	660
45	gacattcttg tcgaacaagg tgcacagcgt cttgtacaag ttggtcttgg agatgatgac	720
	cagtgtattg aagatgactt taccgcttgg cgagaagcat tgtggcccga gcttgataca	780
	atactgaggg aagaagggga tacagctggt gccacacat aactgcagc tgtgttagaa	840
50	tacagagttt ctattcacga ctctgaagat gccaaattca atgatataaa catggcaaat	900
	gggaatggtt aactgtgtt tgatgctcaa catccttaca aagcaaatgt cgctgttaaa	960
55	agggagcttc atactcccgat gtctgatcgt tcttgatatcc atttggaatt tgacattgct	1020
	ggaagtggac ttacgtatga aactggagat catgttggtg tactttgtga taacttaagt	1080
	gaaactgtag atgaagctct tagattgctg gatatgtcac ctgatactta tttctcactt	1140

	cacgctgaaa aagaagacg caccaccaatc agcagctcac tgcctcctcc cttcccacct	1200
	tgcaacttga gaacagcgct tacacgatat gcatgtcttt tgagttctcc aaagaagtct	1260
5	gcttttagttg cgttggctgc tcatgcatct gatcctaccg aagcagaacg attaaaacac	1320
	cttgcttcac ctgctggaaa ggatgaatat tcaaagtggg tagtagagag tcaaagaagt	1380
10	ctacttgagg tgatggccga gtttccttca gccaaagccac cacttggtgt cttcttcgct	1440
	ggagttgctc caaggttgca gcctagggtc tattcgatat catcatcgcc caagattgct	1500
	gaaactagaa ttcacgtcac atgtgcactg gtttatgaga aaatgccaac tggcaggatt	1560
15	cataagggag tgtgttccac ttggatgaag aatgctgtgc cttacgagaa gagtgaaaac	1620
	tgttcctcgg cgccgatatt tgtaggcaa tccaacttca agcttccttc tgattctaag	1680
20	gtaccgatca tcatgatcgg tccagggact ggattagctc cattcagagg attccttcag	1740
	gaaagactag cgttggtaga atctgggtgt gaacttgggc catcagtttt gttctttgga	1800
	tgcagaaacc gtagaatgga tttcatctac gaggaagagc tccagcgatt tgtagagagt	1860
25	ggtgctctcg cagagctaag tgtcgccttc tctcgtgaag gaccaccaa agaatacgta	1920
	cagcacaaga tgatggacaa ggcttctgat atctggaata tgatctctca aggagcttat	1980
30	ttatatgttt gtggtgacgc caaaggcatg gcaagagatg ttcacagatc tctccacaca	2040
	atagctcaag aacaggggtc aatggattca actaaagcag agggcttcgt gaagaatctg	2100
35	caaacgagtg gaagatatct tagagatgta tggtaa	2136
	<210> 18	
	<211> 711	
	<212> PRT	
40	<213> Arabidopsis thaliana	
	<400> 18	
45	Met Ser Ser Ser Ser Ser Ser Ser Thr Ser Met Ile Asp Leu Met Ala	
	1 5 10 15	
	Ala Ile Ile Lys Gly Glu Pro Val Ile Val Ser Asp Pro Ala Asn Ala	
50	20 25 30	
	Ser Ala Tyr Glu Ser Val Ala Ala Glu Leu Ser Ser Met Leu Ile Glu	
	35 40 45	
55	Asn Arg Gln Phe Ala Met Ile Val Thr Thr Ser Ile Ala Val Leu Ile	
	50 55 60	

88

	Gly	Cys	Ile	Val	Met	Leu	Val	Trp	Arg	Arg	Ser	Gly	Ser	Gly	Asn	Ser	
	65					70					75					80	
5	Lys	Arg	Val	Glu	Pro	Leu	Lys	Pro	Leu	Val	Ile	Lys	Pro	Arg	Glu	Glu	
					85					90					95		
10	Glu	Ile	Asp	Asp	Gly	Arg	Lys	Lys	Val	Thr	Ile	Phe	Phe	Gly	Thr	Gln	
				100					105					110			
15	Thr	Gly	Thr	Ala	Glu	Gly	Phe	Ala	Lys	Ala	Leu	Gly	Glu	Glu	Ala	Lys	
			115					120					125				
20	Ala	Arg	Tyr	Glu	Lys	Thr	Arg	Phe	Lys	Ile	Val	Asp	Leu	Asp	Asp	Tyr	
		130					135					140					
25	Ala	Ala	Asp	Asp	Asp	Glu	Tyr	Glu	Glu	Lys	Leu	Lys	Lys	Glu	Asp	Val	
	145					150					155					160	
30	Ala	Phe	Phe	Phe	Leu	Ala	Thr	Tyr	Gly	Asp	Gly	Glu	Pro	Thr	Asp	Asn	
					165					170					175		
35	Ala	Ala	Arg	Phe	Tyr	Lys	Trp	Phe	Thr	Glu	Gly	Asn	Asp	Arg	Gly	Glu	
				180					185					190			
40	Trp	Leu	Lys	Asn	Leu	Lys	Tyr	Gly	Val	Phe	Gly	Leu	Gly	Asn	Arg	Gln	
		195						200					205				
45	Tyr	Glu	His	Phe	Asn	Lys	Val	Ala	Lys	Val	Val	Asp	Asp	Ile	Leu	Val	
		210					215					220					
50	Glu	Gln	Gly	Ala	Gln	Arg	Leu	Val	Gln	Val	Gly	Leu	Gly	Asp	Asp	Asp	
	225					230					235					240	
55	Gln	Cys	Ile	Glu	Asp	Asp	Phe	Thr	Ala	Trp	Arg	Glu	Ala	Leu	Trp	Pro	
					245					250					255		
60	Glu	Leu	Asp	Thr	Ile	Leu	Arg	Glu	Glu	Gly	Asp	Thr	Ala	Val	Ala	Thr	
				260						265				270			
65	Pro	Tyr	Thr	Ala	Ala	Val	Leu	Glu	Tyr	Arg	Val	Ser	Ile	His	Asp	Ser	
			275					280					285				
70	Glu	Asp	Ala	Lys	Phe	Asn	Asp	Ile	Asn	Met	Ala	Asn	Gly	Asn	Gly	Tyr	
		290					295					300					

Thr Val Phe Asp Ala Gln His Pro Tyr Lys Ala Asn Val Ala Val Lys
 305 310 315 320
 5
 Arg Glu Leu His Thr Pro Glu Ser Asp Arg Ser Cys Ile His Leu Glu
 325 330 335
 10
 Phe Asp Ile Ala Gly Ser Gly Leu Thr Tyr Glu Thr Gly Asp His Val
 340 345 350
 15
 Gly Val Leu Cys Asp Asn Leu Ser Glu Thr Val Asp Glu Ala Leu Arg
 355 360 365
 20
 Leu Leu Asp Met Ser Pro Asp Thr Tyr Phe Ser Leu His Ala Glu Lys
 370 375 380
 Glu Asp Gly Thr Pro Ile Ser Ser Ser Leu Pro Pro Pro Phe Pro Pro
 385 390 395 400
 25
 Cys Asn Leu Arg Thr Ala Leu Thr Arg Tyr Ala Cys Leu Leu Ser Ser
 405 410 415
 30
 Pro Lys Lys Ser Ala Leu Val Ala Leu Ala Ala His Ala Ser Asp Pro
 420 425 430
 35
 Thr Glu Ala Glu Arg Leu Lys His Leu Ala Ser Pro Ala Gly Lys Asp
 435 440 445
 40
 Glu Tyr Ser Lys Trp Val Val Glu Ser Gln Arg Ser Leu Leu Glu Val
 450 455 460
 Met Ala Glu Phe Pro Ser Ala Lys Pro Pro Leu Gly Val Phe Phe Ala
 465 470 475 480
 45
 Gly Val Ala Pro Arg Leu Gln Pro Arg Phe Tyr Ser Ile Ser Ser Ser
 485 490 495
 50
 Pro Lys Ile Ala Glu Thr Arg Ile His Val Thr Cys Ala Leu Val Tyr
 500 505 510
 55
 Glu Lys Met Pro Thr Gly Arg Ile His Lys Gly Val Cys Ser Thr Trp
 515 520 525
 Met Lys Asn Ala Val Pro Tyr Glu Lys Ser Glu Asn Cys Ser Ser Ala
 530 535 540

5
 Pro Ile Phe Val Arg Gln Ser Asn Phe Lys Leu Pro Ser Asp Ser Lys
 545 550 555 560
 Val Pro Ile Ile Met Ile Gly Pro Gly Thr Gly Leu Ala Pro Phe Arg
 565 570 575
 10
 Gly Phe Leu Gln Glu Arg Leu Ala Leu Val Glu Ser Gly Val Glu Leu
 580 585 590
 15
 Gly Pro Ser Val Leu Phe Phe Gly Cys Arg Asn Arg Arg Met Asp Phe
 595 600 605
 20
 Ile Tyr Glu Glu Glu Leu Gln Arg Phe Val Glu Ser Gly Ala Leu Ala
 610 615 620
 25
 Glu Leu Ser Val Ala Phe Ser Arg Glu Gly Pro Thr Lys Glu Tyr Val
 625 630 635 640
 30
 Gln His Lys Met Met Asp Lys Ala Ser Asp Ile Trp Asn Met Ile Ser
 645 650 655
 35
 Gln Gly Ala Tyr Leu Tyr Val Cys Gly Asp Ala Lys Gly Met Ala Arg
 660 665 670
 40
 Asp Val His Arg Ser Leu His Thr Ile Ala Gln Glu Gln Gly Ser Met
 675 680 685
 45
 Asp Ser Thr Lys Ala Glu Gly Phe Val Lys Asn Leu Gln Thr Ser Gly
 690 695 700
 Arg Tyr Leu Arg Asp Val Trp
 705 710
 50
 <210> 19
 <211> 33
 <212> DNA
 <213> Arabidopsis thaliana
 <400> 19
 cggaattctc atggatcaaa tcgaagcaat gtt
 55
 <210> 20
 <211> 29
 <212> DNA
 <213> Arabidopsis thaliana

	<400> 20	
	cgactagttt agcaaatcgg aatcggagc	29
5		
	<210> 21	
	<211> 30	
	<212> DNA	
	<213> Arabidopsis thaliana	
10		
	<400> 21	
	cgctcgagat atggacctcc tcttgctgga	30
15		
	<210> 22	
	<211> 32	
	<212> DNA	
	<213> Arabidopsis thaliana	
20		
	<400> 22	
	cggggtacctt aacagttcct tggtttcata ac	32
25		
	<210> 23	
	<211> 36	
	<212> DNA	
	<213> Arabidopsis thaliana	
30		
	<400> 23	
	gctctagacc tatggcgcca caagaacaag cagttt	36
35		
	<210> 24	
	<211> 36	
	<212> DNA	
	<213> Arabidopsis thaliana	
40		
	<400> 24	
	gcggatcccc ttcacaatcc atttgctagt tttgcc	36
45		
	<210> 25	
	<211> 31	
	<212> DNA	
	<213> Rheum tataricum	
50		
	<400> 25	
	ccggatccaa atggccccag aagagagcag g	31
55		
	<210> 26	
	<211> 35	
	<212> DNA	
	<213> Rheum tataricum	
	<400> 26	
	cgctcgagtt aagtgatcaa tggaaccgaa gacag	35

	<210> 27	
	<211> 38	
	<212> DNA	
	<213> Rhodobacter capsulatus	
5	<400> 27	
	ccgaattccc atgaccctgc aatctcaaac agctaaag	38
10	<210> 28	
	<211> 30	
	<212> DNA	
	<213> Rhodobacter capsulatus	
15	<400> 28	
	ccactagttt aagcaggtgg atcggcagct	30
20	<210> 29	
	<211> 37	
	<212> DNA	
	<213> Saccharomyces cerevisiae	
25	<400> 29	
	ccctcgagat catgccgttt ggaatagaca acaccga	37
30	<210> 30	
	<211> 36	
	<212> DNA	
	<213> Saccharomyces cerevisiae	
35	<400> 30	
	ccaagcttat cgggctgatt accagacatc ttcttg	36
40	<210> 31	
	<211> 36	
	<212> DNA	
	<213> Arabidopsis thaliana	
45	<400> 31	
	ccggatcccc atgtcctctt cttcttcttc gtcaac	36
50	<210> 32	
	<211> 34	
	<212> DNA	
	<213> Arabidopsis thaliana	
55	<400> 32	
	ccctcgaggt gagtgtgtgg cttcaatagt ttcg	34
55	<210> 33	
	<211> 40	
	<212> DNA	
	<213> Rhodobacter capsulatus	

	<400> 33	
	ccgctcgagc ggatgaccct gcaatctcaa acagctaaag	40
5	<210> 34	
	<211> 30	
	<212> DNA	
	<213> Rhodobacter capsulatus	
10	<400> 34	
	gcggatcctt aagcaggtgg atcggcagct	30
15	<210> 35	
	<211> 35	
	<212> DNA	
	<213> Arabidopsis thaliana	
20	<400> 35	
	tgccatggca atggcgccac aagaacaagc agttt	35
25	<210> 36	
	<211> 36	
	<212> DNA	
	<213> Arabidopsis thaliana	
30	<400> 36	
	gcggatcccc ttcacaatcc atttgctagt ttgccc	36
35	<210> 37	
	<211> 37	
	<212> DNA	
	<213> Arabidopsis thaliana	
40	<400> 37	
	ttgcggccgc aaatctcgat cccgcgaaat taatagc	37
45	<210> 38	
	<211> 36	
	<212> DNA	
	<213> Arabidopsis thaliana	
50	<400> 38	
	cgctcgagcc ttcacaatcc atttgctagt ttgccc	36
55		

Claims

- 5 1. A micro-organism having an operative metabolic pathway comprising at least one enzyme activity, said pathway producing 4-coumaric acid and producing resveratrol therefrom, or an oligomeric or glycosidically-bound derivative thereof.
- 10 2. A micro-organism as claimed in claim 1, wherein said resveratrol is produced in a reaction catalysed by an enzyme in which endogenous malonyl-CoA is a substrate.
- 15 3. A micro-organism as claimed in claim 1 or claim 2, wherein said resveratrol is produced from 4-coumaroyl-CoA.
- 20 4. A micro-organism as claimed in claim 3, wherein said resveratrol is produced from 4-coumaroyl-CoA by a resveratrol synthase.
- 25 5. A micro-organism as claimed in claim 4, wherein said resveratrol synthase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.
- 30 6. A micro-organism as claimed in claim 5, wherein said resveratrol synthase is resveratrol synthase (EC 2.3.1.95) from a plant belonging to the genus of *Arachis*, a plant belonging to the genus of *Rheum*, or a plant belonging to the genus of *Vitus* or any one of the genera *Pinus*, *Picea*, *Lilium*, *Eucalyptus*, *Parthenocissus*, *Cissus*, *Calochortus*,

Polygonum, Gnetum, Artocarpus, Nothofagus, Phoenix, Festuca, Carex, Veratrum, Bauhinia or Pterolobium.

7. A micro-organism as claimed in any preceding claim,
5 wherein said 4-coumaric acid is produced from *trans*-cinnamic acid.
8. A micro-organism as claimed in claim 7, wherein said
4-coumaric acid is produced from *trans*-cinnamic acid by an
10 enzyme in a reaction catalysed by said enzyme in which oxygen is a substrate, NADH or NADPH is a cofactor and NAD or NADP is a product.
9. A micro-organism as claimed in claim 8, wherein said
15 4-coumaric acid is produced from *trans*-cinnamic acid by a cinnamate 4-hydroxylase.
10. A micro-organism as claimed in claim 9, wherein said
cinnamate 4-hydroxylase is expressed in said micro-
20 organism from nucleic acid coding for said enzyme which is not native to the micro-organism.
11. A micro-organism as claimed in claim 10, wherein said
cinnamate 4-hydroxylase is cinnamate 4-hydroxylase (EC
25 1.14.13.11) from a plant belonging to the genus of *Arabidopsis*, a plant belonging to the genus of *Citrus*, a plant belonging to the genus of *Phaseolus*, a plant belonging to the genus of *Pinus*, a plant belonging to the genus of *Populus*, a plant belonging to the genus of
30 *Solanum*, a plant belonging to the genus of *Vitus*, a plant belonging to the genus of *Zea*, or any one of the genera *Ammi*, *Avicennia*, *Camellia*, *Camptotheca*, *Catharanthus*, *Glycine*, *Helianthus*, *Lotus*, *Mesembryanthemum*,

Physcomitrella, *Ruta*, *Saccharum*, and *Vigna*, from a filamentous fungus belonging to the genus *Aspergillus*.

12. A micro-organism as claimed in any one of claims 1 to
5 6, wherein said 4-coumaric acid is produced from tyrosine in a reaction catalysed by an enzyme in which ammonia is produced.
13. A micro-organism as claimed in any one of claims 1 to
10 4, wherein said 4-coumaric acid is produced from *tyrosine* by a L-phenylalanine ammonia lyase or a tyrosine ammonia lyase.
14. A micro-organism as claimed in claim 13, wherein said
15 4-coumaric acid is produced by tyrosine ammonia lyase (EC 4.3.1.5) from yeast or bacteria.
15. A micro-organism as claimed in claim 14, wherein said
20 tyrosine ammonia lyase is from the yeast *Rhodotorula rubra* or from the bacterium *Rhodobacter capsulatus*.
16. A micro-organism as claimed in any one of claims 12 to
25 15, wherein said tyrosine ammonia lyase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.
17. A micro-organism as claimed in any one of claims 9 to
30 11, wherein said *trans*-cinnamic acid is produced from L-phenylalanine in a reaction catalysed by an enzyme in which ammonia is produced.

18. A micro-organism as claimed in claim 17, wherein said *trans*-cinnamic acid is formed from L-phenylalanine by a L-phenylalanine ammonia lyase.
- 5 19. A micro-organism as claimed in claim 18, wherein said *trans*-cinnamic acid is formed from L-phenylalanine by L-phenylalanine ammonia lyase (EC 4.3.1.5) from a plant
- 10 belonging to the genus of *Arabidopsis*, a plant belonging to the genus of *Brassica*, a plant belonging to the genus of *Citrus*, a plant belonging to the genus of *Phaseolus*, a plant belonging to the genus of *Pinus*, a plant belonging to the genus of *Populus*, a plant belonging to the genus of *Solanum*, a plant belonging to the genus of *Prunus*, a plant belonging to the genus of *Vitus*, a plant belonging to the
- 15 genus of *Zea*, or a plant belonging to any one of the genera *Agastache*, *Ananas*, *Asparagus*, *Bromheadia*, *Bambusa*, *Beta*, *Betula*, *Cucumis*, *Camellia*, *Capsicum*, *Cassia*, *Catharanthus*, *Cicer*, *Citrullus*, *Coffea*, *Cucurbita*, *Cynodon*, *Daucus*, *Dendrobium*, *Dianthus*, *Digitalis*, *Dioscorea*, *Eucalyptus*, *Gallus*, *Ginkgo*, *Glycine*, *Hordeum*, *Helianthus*, *Ipomoea*, *Lactuca*, *Lithospermum*, *Lotus*, *Lycopersicon*, *Medicago*, *Malus*, *Manihot*, *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Olea*, *Oryza*, *Pisum*, *Persea*, *Petroselinum*, *Phalaenopsis*, *Phyllostachys*, *Physcomitrella*,
- 20 *Picea*, *Pyrus*, *Quercus*, *Raphanus*, *Rehmannia*, *Rubus*, *Sorghum*, *Sphenostylis*, *Stellaria*, *Stylosanthes*, *Triticum*, *Trifolium*, *Triticum*, *Vaccinium*, *Vigna*, or *Zinnia*, or from a filamentous fungus belonging to the genus *Aspergillus*.
- 25
- 30 20. A micro-organism as claimed in any one of claims 18 or claim 19, wherein said phenylalanine ammonia lyase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.

21. A micro-organism as claimed in any preceding claim,
wherein 4-coumaroyl-CoA is formed in a reaction catalysed
by an enzyme in which ATP and CoA are substrates and ADP
5 is a product.
22. A micro-organism as claimed in any preceding claim,
wherein 4-coumaroyl-CoA is formed in a reaction catalysed
by a 4-coumarate-CoA ligase.
- 10 23. A micro-organism as claimed in claim 22, wherein said
4-coumarate-CoA ligase is 4-coumarate-CoA ligase (EC
6.2.1.12) from a plant belonging to the genus of *Abies*, a
plant belonging to the genus of *Arabidopsis*, a plant
15 belonging to the genus of *Brassica*, a plant belonging to
the genus of *Citrus*, a plant belonging to the genus of
Larix, a plant belonging to the genus of *Phaseolus*, a
plant belonging to the genus of *Pinus*, a plant belonging
to the genus of *Populus*, a plant belonging to the genus of
20 *Solanum*, a plant belonging to the genus of *Vitus*, a plant
belonging to the genus of *Zea*, or a plant belonging to any
one of the genera *Agastache*, *Amorpha*, *Cathaya*, *Cedrus*,
Crocus, *Festuca*, *Glycine*, *Juglans*, *Keteleeria*,
Lithospermum, *Lolium*, *Lotus*, *Lycopersicon*, *Malus*,
25 *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Nothotsuga*, *Oryza*,
Pelargonium, *Petroselinum*, *Physcomitrella*, *Picea*, *Prunus*,
Pseudolarix, *Pseudotsuga*, *Rosa*, *Rubus*, *Ryza*, *Saccharum*,
Suaeda, *Thellungiella*, *Triticum*, or *Tsuga*, from a
filamentous fungus belonging to the genus *Aspergillus*, a
30 filamentous fungus belonging to the genus *Neurospora*, a
fungus belonging to the genus *Yarrowia*, a fungus belonging
to the genus of *Mycosphaerella*, from a bacterium belonging
to the genus of *Mycobacterium*, a bacterium belonging to

the genus of *Neisseria*, a bacterium belonging to the genus of *Streptomyces*, a bacterium belonging to the genus of *Rhodobacter*, or from a nematode belonging to the genus *Ancylostoma*, a nematode belonging to the genus *Caenorhabditis*, a nematode belonging to the genus *Haemonchus*, a nematode belonging to the genus *Lumbricus*, a nematode belonging to the genus *Meilodogyne*, a nematode belonging to the genus *Strongyloidus*, or a nematode belonging to the genus *Pristionchus*.

10

24. A micro-organism as claimed in any preceding claim, wherein at least one copy of at least one genetic sequence encoding a respective enzyme in said metabolic pathway has been recombinantly introduced into said micro-organism.

15

25. A micro-organism as claimed in claim 24, wherein a native NADPH:cytochrome P450 reductase (CPR) has been overexpressed in said micro-organism.

20 26. A micro-organism as claimed in claim 24, wherein a NADPH:cytochrome P450 reductase (CPR) has been recombinantly introduced into said micro-organism.

25 27. A micro-organism as claimed in claim 26, wherein said NADPH:cytochrome P450 reductase is NADPH:cytochrome P450 reductase (EC 1.6.2.4) from a plant belonging to the genus of *Arabidopsis*, e.g. *A. thaliana*, a plant belonging to the genus of *Citrus*, e.g. *C. sinensis*, *C. x paradisi*, a plant belonging to the genus of *Phaseolus*, e.g. *P. vulgaris*, a plant belonging to the genus of *Pinus*, e.g. *P. taeda*, a plant belonging to the genus of *Populus*, e.g. *P. deltoides*, *P. tremuloides*, *P. trichocarpa*, a plant belonging to the genus of *Solanum*, e.g. *S. tuberosum*, a

30

plant belonging to the genus of *Vitus*, e.g. *Vitus vinifera*, a plant belonging to the genus of *Zea*, e.g. *Z. mays*, or other plant genera e.g. *Ammi*, *Avicennia*, *Camellia*, *Camptotheca*, *Catharanthus*, *Glycine*, *Helianthus*,
5 *Lotus*, *Mesembryanthemum*, *Physcomitrella*, *Ruta*, *Saccharum*, *Vigna*.

28. A micro-organism as claimed in any preceding claim,
wherein at least one copy of a genetic sequence encoding a
10 tyrosine ammonia lyase is operatively linked to an
expression signal not natively associated with said
genetic sequence in said organism.

29. A micro-organism as claimed in any preceding claim
15 wherein at least one copy of a genetic sequence encoding a
phenylalanine ammonia lyase is operatively linked to an
expression signal not natively associated with said
genetic sequence in said organism.

20 30. A micro-organism as claimed in any preceding claim
wherein at least one copy of a genetic sequence encoding
cinnamate 4-hydroxylase is operatively linked to an
expression signal not natively associated with said
genetic sequence in said organism.

25
31. A micro-organism as claimed in any preceding claim
wherein at least one copy of a genetic sequence encoding a
4-coumarate-CoA ligase is operatively linked to an
expression signal not natively associated with said
30 genetic sequence in said organism.

32. A micro-organism as claimed in any preceding claim
wherein at least one copy of a genetic sequence encoding a

resveratrol synthase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

5 33. A micro-organism as claimed in any preceding claim, which is a fungus.

34. A micro-organism as claimed in claim 33, which is a filamentous fungi.

10

35. A micro-organism as claimed in claim 34, which is a micro-organism belonging to the genus *Aspergillus*.

15 36. A micro-organism as claimed in claim 35, which is a strain of *Aspergillus niger* or *A. oryzae*.

37. A micro-organism as claimed in 33, which is a yeast.

20 38. A micro-organism as claimed in claim 37, which is a micro-organism belonging to the genus *Saccharomyces*, *Kluveromyces*, *Candida*, *Pichia*, *Debaromyces*, *Hansenula*, *Yarrowia*, *Zygosaccharomyces* or *Schizosaccharomyces*.

25 39. A micro-organism as claimed in claim 38, which is a strain of *Saccharomyces cerevisiae*, *S. kluyveri*, *S. bayanus*, *S. exiguus*, *S. sevazzi*, *S. uvarum*, *Kluveromyces lactis*, *K. marxianus* var. *marxianus*, *K. thermotolerans*, *Candida utilis*, *C. tropicalis*, *Pichia stipidis*, *P. pastoris*, *P. sorbitophila*, *Debaromyces hansenii*, *Hansenula polymorpha*, *Yarrowia lipolytica*, *Zygosaccharomyces rouxii*
30 or *Schizosaccharomyces pombe*.

40. A micro-organism as claimed in any one of claim 1 to 32, which is a bacterium.
41. A micro-organism as claimed in claim 40, which is a
5 micro-organism belonging to the genus *Escherichia* or *Lactococcus*.
42. A micro-organism as claimed in claim 41, which is a strain of *Escherichia coli* or *Lactococcus lactis*.
10
43. A micro-organism as claimed in claim 1, containing one or more copies of an heterologous DNA sequence encoding phenylalanine ammonia lyase operatively associated with an expression signal, and containing one or more copies of an
15 heterologous DNA sequence encoding cinnamate-4-hydroxylase operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding 4-coumarate CoA-ligase operatively associated with an expression signal, and containing one
20 or more copies of an heterologous DNA sequence encoding resveratrol synthase operatively associated with an expression signal.
44. A micro-organism as claimed in claim 1, lacking
25 cinnamate-4-hydroxylase activity, and containing one or more copies of a heterologous DNA sequence encoding tyrosine ammonia lyase operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding 4-coumarate CoA-ligase
30 operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding resveratrol synthase operatively associated with an expression signal.

45. A method for producing resveratrol or an oligomeric or glycosidically-bound derivative thereof comprising contacting a non-plant cell having an appropriate
5 resveratrol producing metabolic pathway with a carbon substrate in the substantial absence of an external source of 4-coumaric acid.
46. A method as claimed in claim 45, wherein said non-
10 plant cell is selected from the group consisting of fungi and bacteria.
47. A method as claimed in claim 46, where said non-plant
15 cell is a fungus selected from the group of yeast.
48. A method as claimed in claim 47, where said yeast is
selected from the species *Saccharomyes*.
49. A method as claimed in any one of claims 45 to 48,
20 wherein said non-plant cell is of a micro-organism as claimed in any one of claims 1 to 44.
50. A method as claimed in any one of claim 45 to 49,
25 wherein said carbon substrate is selected from the group of fermentable carbon substrates consisting of monosaccharides, oligosaccharides and polysaccharides.
51. A method as claimed in claim 50, wherein said
30 fermentable carbon substrate is glucose, fructose, galactose, xylose, arabinose, mannose, sucrose, lactose, erythrose, threose, ribose.

52. A method as claimed in any one of claim 45 to 49, wherein said carbon substrate is selected from the group of non-fermentable carbon substrate.
- 5 53. A method as claimed in claim 52, wherein said non-fermentable carbon substrate is ethanol, acetate, glycerol, lactate.
54. A method as claimed in claim 52, wherein said non-fermentable carbon substrate is selected from the group consisting of amino acids.
- 10 55. A method as claimed in claim 54, wherein said non-fermentable carbon substrate is selected from the group consisting of phenylalanine and tyrosine.
- 15 56. A method for producing resveratrol or an oligomeric or glycosidically-bound derivative thereof through heterologous expression of nucleotide sequences encoding phenylalanine ammonia lyase, cinnamate 4-hydroxylase, 4-coumarate-CoA ligase and resveratrol synthase.
- 20 57. A method for producing resveratrol or an oligomeric or glycosidically-bound derivative thereof through heterologous expression of nucleotide sequences encoding tyrosine ammonia lyase, 4-coumarate-CoA ligase and resveratrol synthase.
- 25 58. A method as claimed in any one of claims 45 to 57, further including using said produced resveratrol or an oligomeric or glycosidically-bound derivative thereof as a nutraceutical in a dairy product or a beverage.
- 30

59. A method as claimed in claim 58, wherein said resveratrol or an oligomeric or glycosidically-bound derivative thereof is used as a nutraceutical in beer.

1/7




Figure 1

2/7




Figure 2

3/7




Figure 3

4/7





Figure 4

5/7




uv-spectrum of pure
trans-resveratrol
60 nanogram total

uv-spectrum of *trans*-resveratrol
in extract of *S. cerevisiae* strain
FSSC-PALC4H4CLVST
(PAL-pathway)

uv-spectrum of *trans*-resveratrol
in extract of *S. cerevisiae* strain
FSSC-TAL4CLVST
(TAL-pathway)

Figure 5

6/7



E. coli strain:
FSEC-control
(empty vectors)

E. coli strain:
FSEC-TAL4CLVST
(TAL-pathway)

Figure 6

7/7




Figure 7

INTERNATIONAL SEARCH REPORT

 International application No
 PCT/EP2006/060154

A. CLASSIFICATION OF SUBJECT MATTER

INV. C12N9/00 C12N9/02 C12N9/10 C12N9/88 C12P7/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)
 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SAMAPPITO S ET AL: "Aromatic and pyrone polyketides synthesized by a stilbene synthase from Rheum tataricum" PHYTOCHEMISTRY, PERGAMON PRESS, GB, vol. 62, no. 3, February 2003 (2003-02), pages 313-323, XP004412246 ISSN: 0031-9422 cited in the application abstract ----- -/--	45,46



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

12 June 2006

Date of mailing of the international search report

20/06/2006

Name and mailing address of the ISA/

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Huse, I

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2006/060154

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BECKER JOHN V W ET AL: "Metabolic engineering of <i>Saccharomyces cerevisiae</i> for the synthesis of the wine-related antioxidant resveratrol." FEMS YEAST RESEARCH. OCT 2003, vol. 4, no. 1, October 2003 (2003-10), pages 79-85, XP002347057 ISSN: 1567-1356 cited in the application the whole document	1-59
Y	----- WATTS KEVIN T ET AL: "Exploring recombinant flavonoid biosynthesis in metabolically engineered <i>Escherichia coli</i> ." CHEMBIOCHEM : A EUROPEAN JOURNAL OF CHEMICAL BIOLOGY. 2 APR 2004, vol. 5, no. 4, 2 April 2004 (2004-04-02), pages 500-507, XP002347112 ISSN: 1439-4227 cited in the application abstract	1-59
Y	----- HWANG E I ET AL: "Production of plant-specific flavanones by <i>Escherichia coli</i> containing an artificial gene cluster" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, WASHINGTON, DC, US, vol. 69, no. 5, May 2003 (2003-05), pages 2699-2706, XP002977268 ISSN: 0099-2240 cited in the application abstract the whole document	1-59
A	----- RO DAE-KYUN ET AL: "Reconstitution of the entry point of plant phenylpropanoid metabolism in yeast (<i>Saccharomyces cerevisiae</i>): implications for control of metabolic flux into the phenylpropanoid pathway." THE JOURNAL OF BIOLOGICAL CHEMISTRY. 23 JAN 2004, vol. 279, no. 4, 23 January 2004 (2004-01-23), pages 2600-2607, XP002347058 ISSN: 0021-9258 cited in the application abstract page 2602, right-hand column - page 2603, left-hand column ----- -/--	

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2006/060154

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02/10407 A (E.I. DUPONT DE NEMOURS AND COMPANY) 7 February 2002 (2002-02-07) abstract page 13, line 12 - line 36 -----	
A	JEANDET PHILIPPE ET AL: "Phytoalexins from the Vitaceae: biosynthesis, phytoalexin gene expression in transgenic plants, antifungal activity, and metabolism." JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY. 8 MAY 2002, vol. 50, no. 10, 8 May 2002 (2002-05-08), pages 2731-2741, XP002347113 ISSN: 0021-8561 the whole document -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2006/060154

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0210407 A	07-02-2002	EP 1303622 A1 JP 2004530402 T	23-04-2003 07-10-2004